

Correction

Open Access

Trace metals and over-expression of metallothioneins in bladder tumoral lesions: a case-control study

André FS Amaral *^{1,2,3}, Teresa Cymbron^{2,4}, Fátima Gärtner⁵, Manuela Lima^{2,4} and Armindo S Rodrigues^{2,3}

Address: ¹Spanish National Cancer Research Centre (CNIO), Genetic and Molecular Epidemiology Group, Madrid, Spain, ²CIRN, DB, University of the Azores, APT 1422, 9501-855 Ponta Delgada, Portugal, ³PHERG, University of the Azores, APT 1422, 9501-855 Ponta Delgada, Portugal,

⁴IBMC-Institute for Molecular and Cell Biology, University of Porto, 4150-180 Porto, Portugal and ⁵IPATIMUP, R. Dr. Roberto Frias, s/n, University of Porto, 4200-456 Porto, Portugal

Email: André FS Amaral* - aamaral@cnio.es; Teresa Cymbron - tcymbron@uac.pt; Fátima Gärtner - fgartner@ipatimup.pt; Manuela Lima - mlima@uac.pt; Armindo S Rodrigues - rodrigues@uac.pt

* Corresponding author

Published: 11 November 2009

BMC Veterinary Research 2009, 5:40 doi:10.1186/1746-6148-5-40

This article is available from: <http://www.biomedcentral.com/1746-6148/5/40>

© 2009 Amaral et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Received: 30 October 2009

Accepted: 11 November 2009

Correction

During the publication process of this work [1], we noted that the column for nickel in table 2 was not showing the concentration values in the proper units. This change has been made in the updated table (see table 2).

References

- I. Amaral AFS, Cymbron T, Gartner F, Lima M, Rodrigues AS: **Trace metals and over-expression of metallothioneins in bladder tumoral lesions: a case-control study.** *BMC Veterinary Research* 2009, **5**:24.

Table 2: Concentrations (mean $\mu\text{g} \cdot \text{kg}^{-1} \pm$ standard error) of arsenic, cadmium, chromium, lead and nickel as measured in hair and bladder tissue, from controls and case lesions.

	Arsenic	Cadmium	Chromium	Lead	Nickel
Bladder tissue					
Controls (n = 17)	19.94 ± 6.46	21.60 ± 2.02	269.41 ± 38.18	10.29 ± 1.84	80 ± 30
All cases (n = 37)	23.00 ± 2.62	25.91 ± 1.95	247.84 ± 33.36	14.73 ± 2.87	70 ± 10
Carcinomas (n = 18)	17.94 ± 4.21	20.83 ± 2.13	283.89 ± 41.85	13.33 ± 3.77	60 ± 10
Hair					
Controls (n = 13)	59.54 ± 12.44	16.25 ± 3.01	123.85 ± 25.83	460.77 ± 176.78	90 ± 20
All cases (n = 23)	61.96 ± 8.25	23.20 ± 2.39 ^a	634.35 ± 127.78 ^a	1095.22 ± 214.77 ^a	300 ± 50 ^a
Carcinomas (n = 10)	58.30 ± 12.69	21.61 ± 3.44	559.00 ± 198.40 ^a	960.00 ± 318.04	280 ± 80 ^a

^a differs significantly ($p \leq 0.05$) from controls.