

RESEARCH ARTICLE

Open Access

# The increase of methicillin-resistant *Staphylococcus aureus* (MRSA) and the presence of an unusual sequence type ST49 in slaughter pigs in Switzerland

Gudrun Overesch<sup>1,2</sup>, Sabina Büttner<sup>3</sup>, Alexandra Rossano<sup>1</sup> and Vincent Perreten<sup>1\*</sup>

## Abstract

**Background:** In years past, methicillin-resistant *S. aureus* (MRSA) has been frequently detected in pigs in Europe, North America and Asia. Recent, yet sporadic studies have revealed a low occurrence of MRSA in Switzerland. In 2009, a monitoring survey of the prevalence and genetic diversity of methicillin-resistant *S. aureus* (MRSA) in slaughter pigs in Switzerland was conducted using methods recommended by the EU guidelines, and using a sampling strategy evenly distributed throughout the year and representative of the Swiss slaughter pig population. Monitoring should determine if the overall prevalence of MRSA in the entire country is increasing over the years and if specific multi-resistant MRSA clones are spreading over the country.

**Results:** In 2009, the nasal cavities of eight out of 405 randomly selected pigs were positive for MRSA, representing a prevalence of 2.0% (95% CI 0.9-3.9). The following year, 23 out of 392 pigs were positive for MRSA [5.9% prevalence (95% CI 3.8-8.7)]. Three multilocus sequence types (ST), four *spa* types and two types of staphylococcal cassette chromosome *mec* (SCC*mec*) elements were detected. The most frequent genotypes were ST398 (MLST)-(*spa*)t034-V (SCC*mec*) (n = 18) and ST49-t208-V (n = 7), followed by ST398-t011-V (n = 4), ST398-t1451-V (n = 1), and ST1-t2279-IVc (n = 1). The isolates displayed resistance to  $\beta$ -lactams [*mecA*, (31/31); *blaZ*, (19/31)]; tetracycline [*tet(M)*, (31/31); *tet(K)*, (30/31)] (n = 31); macrolides and lincosamides [*erm(C)* (4/31) or *erm(A)* (18/31)] (n = 22); tiamulin [*vga(A)v* (9/31) or unknown mechanism (18/31)] (n = 27); trimethoprim [*dhfr(G)* (18/31)]; spectinomycin [*ant(9)-Ia* (19/31) or unknown mechanism (3/31)] (n = 22); streptomycin [*str* (19/31)]; sulphamethoxazole (7/31) and ciprofloxacin (n = 1) (mechanisms not determined).

**Conclusions:** This study is the first to describe the presence of MRSA ST49 in slaughter pigs, and to demonstrate a significant and nearly three-fold increase of MRSA prevalence in pigs within two years. The presence of a specific clonal lineage of MRSA from Switzerland suggests that it has been selected in Swiss pig husbandry. Effective hygiene measures should be enhanced within the entire pig production chain to suppress the spread of these pathogens into the community.

## Background

Over the years, methicillin-resistant *Staphylococcus aureus* (MRSA) has been increasingly reported in animals worldwide [1-4]. A specific clonal lineage, MLST ST398, has been found to be widespread among pigs in Europe and North America, whereas ST9 seemed to be predominant

in Asia [4-6]. Several *spa* types were detected within these lineages with *spa* types t011, t108, and t034 being the most frequent in MRSA ST398, and t899 in MRSA ST9 [3-6]. Colonisation with MRSA ST398 was also described in other animals like poultry, horses, and calves, and recently, ST398 has been associated with infections in animals and humans [2,4,7]. Humans that come in close contact with farm animals were more likely to be colonised with livestock-associated ST398 and had a higher risk of developing infections with ST398 in case of hospitalisation

\* Correspondence: vincent.perreten@vetsuisse.unibe.ch

<sup>1</sup>Institute of Veterinary Bacteriology, Vetsuisse Faculty, University of Bern, Länggass-Strasse 122, CH-3012 Bern, Switzerland

Full list of author information is available at the end of the article

[8-18]. The pig husbandry environment also represents a large reservoir for MRSA. MRSA ST398 was recovered from dust samples in pig production holdings in Europe with an average prevalence of 26.9% varying between 0% and 51.2% among the EU member states in 2008 [19]. MRSA ST398-t011 was the most dominant type (63.5%), followed by ST398-t108 (9.4%), and ST398-t034 (8.4%). 1.4% of the MRSA from this study were from seven other non-ST398 strains (ST1, ST5, ST8, ST9, ST39, ST97, and ST132). The majority of the non-ST398 strains originated from Germany and Italy [19].

In 2008, dust samples from pig husbandries in Switzerland were analysed and did not contain MRSA [19]. In fact, there was no MRSA detected during previous studies in Swiss slaughter pigs and pig carcasses before 2009 [20,21]. However in 2009, 10 MRSA ST398-t034, which contained the staphylococcal cassette chromosome *SCCmecV* (ST398-t034-V), were found among 800 nasal swabs from Swiss pigs at two slaughterhouses; these results indicated that MRSA had also emerged in the Swiss pig population [22]. That same year, an official monitoring of MRSA in pigs at the slaughterhouse was launched to determine the overall prevalence in the entire country. A sampling strategy for monitoring antimicrobial resistance was developed in 2008, which considered the geographical distribution and size of the slaughterhouses, and the number of slaughtered pigs in Switzerland [23]. Since 2009, this sampling plan was also used for monitoring MRSA. This study will determine the overall prevalence and dynamics of MRSA colonisation in the slaughtered pig population and will indicate if specific genetic lineages of MRSA are colonising pigs in Switzerland.

## Results

### Increased prevalence and molecular typing

MRSA (n = 31) were detected in samples from 5 of 9 slaughterhouses (A, B, C, E, G) in 2009 and in 7 of 9 slaughterhouses (A-D, G-I) in 2010 (Table 1). MRSA was mostly found in pigs raised in cantons where the pig population is the highest (Figure 1), that is, Thurgau (n = 11), Lucerne (n = 6), St. Gallen (n = 5), Bern (n = 4), and Aargau (n = 2). Single isolates were found in pigs from Appenzell Innen Rhodes, Jura, and Zurich (Table 1 and Table 2). In 2009, the prevalence of MRSA in Swiss slaughter pigs was 2.0% (95% CI 0.9-3.9) with eight out of 405 pig nasal samples being positive and increased significantly (p-value < 0.05) to 5.9% (95% CI 3.8-8.7) in 2010 with 23 out of 392 nasal swabs containing MRSA. MRSA was detected every month in 2010, except in September, with up to five positive samples per month, whereas MRSA was only sporadically detected over the year in 2009.

With the exception of one isolate (*SCCmecIVc*), most of the MRSA contained the *SCCmecV* element (Table 2).

The eight MRSA isolated in 2009 belonged to five genotypes: ST398-t011-V (n = 3), ST398-t034-V (n = 1), ST398-t1451-V (n = 1), ST49-t208-V (n = 2), and ST1-t2279-IVc (n = 1); each isolate displayed a different resistance profile (Table 2). In 2010, only three clonal lineages were identified with ST398-t034-V (n = 17) and ST49-t208-V (n = 5) being the most common, whereas ST398-t011-V was only detected once (Table 1 and Table 2).

MRSA isolated from a same slaughterhouse in 2009 and 2010 displayed a different genetic profile and were isolated at different days, except for some MRSA ST398-t034 and ST49-t208 isolates. They were also detected several times from samples from the same slaughterhouses at different days [slaughterhouses A, C, D (ST398-t034)] as well as during the same sampling day in 2010 [slaughterhouses A (ST398-t034) and G (ST49-t208)]. All samples originated from pigs raised in different husbandry (Table 1 and Table 2).

### Resistance profiles

Isolates belonging to the most commonly detected genotype, ST398-t034-V, shared the same resistance profile; however, one isolate was susceptible to streptomycin and did not contain the streptomycin adenylyltransferase gene *str* (Table 2). Otherwise, ST398-t034-V isolates showed resistance to  $\beta$ -lactams specified by *mecA* and *blaZ*, tetracycline [*tet(K)*, *tet(M)*], macrolide-lincosamide-streptogramin B (MLS<sub>B</sub>) antibiotics [*erm(A)*], spectinomycin [*ant(9)-Ia*], streptomycin [*str*], trimethoprim [*dhfr(G)*], and tiamulin. The tiamulin resistance could not be attributed to either of the five known pleuromutilin-streptogramin A resistance genes *vga(A)*, *vga(A)v*, *vga(B)*, *vga(C)*, *vga(D)* [24], or to the *cfp* rRNAmethylase transferase gene which confers cross resistance to phenicols, lincosamides, oxazolidones, pleuromutilins, and streptogramin A antibiotics [25]. Mutations in the 23 rRNA and *rplC* gene were not investigated [26].


Different resistance profiles were observed among the ST49-t208-V isolates, which was the second most prevalent lineage. The difference was mainly due to the presence, or absence, of either of the MLS<sub>B</sub> resistance genes *erm(A)* or *erm(C)* (Table 2). In general, ST49 isolates displayed resistance to  $\beta$ -lactams [*mecA*], tetracycline [*tet(K)*, *tet(M)*], tiamulin [*vga(A)v*], and sulphonamides (these mechanisms were not characterised). Notably, ST49-t208 isolates were susceptible to trimethoprim and resistant to sulphonamides, whereas ST398-t034 isolates were resistant to trimethoprim and susceptible to sulphonamides.

Different antibiotic resistance profiles were also found among the less frequent isolates, ST398-t011 and ST398-t1451. They all displayed resistance to  $\beta$ -lactams [*mecA*] and tetracycline [*tet(K)*, *tet(M)*]; however, resistance to MLS<sub>B</sub> antibiotics [*erm(C)*], tiamulin [*vga(A)v*], streptomycin [*str*], and spectinomycin varied from strain

**Table 1 Distribution of methicillin-resistant *Staphylococcus aureus* genotypes in slaughterhouses and pig farms in Switzerland**

Abattoir	2009					2010				
	Total no. of samples	No. of MRSA positive samples	Percentage of MRSA positive samples	Origin <sup>a</sup>	Genotype (n)	Total no. of samples	No. of MRSA positive samples	Percentage of MRSA positive samples	Origin <sup>a</sup>	Genotype (n)
A	84	2	2.4	TG1, LU2	ST398-t011-V (2)	68	11	16.2	LU4, LU5, SG3, SG5, TG5, TG8, TG9, BE3, BE4 ZH1, JU1	ST398-t034-V (11)
B	90	1	1.1	SG1	ST398-t011-V	101	2	2.0	AG2, SG6	ST49-t208-V (2)
C	63	1	1.6	BE1	ST398-t034-V	65	4	6.2	TG6 BE2, TG7, LU6	ST49-t208-V ST398-t034-V (3)
D	62	0	0	-	-	58	2	3.4	AG1, TG10	ST398-t034-V (2)
E	39	1	2.6	LU1	ST398-t1451-V	37	0	0	-	-
F	28	0	0	-	-	22	0	0	-	-
G	19	3	15.8	TG3, TG2, SG2	ST1-t2279-IVc ST49-t208-V (2)	15	2	13.3	TG4, SG4	ST49-t208-V (2)
H	10	0	0	-	-	14	1	7.1	LU3	ST398-t034-V
I	10	0	0	-	-	12	1	8.3	AI1	ST398-t011-V
A-I (all)	405	8	2.0			392	23	5.9		

a) Geographical origin of the farms: letters indicate the Cantons (AI, Appenzell Innen Rhodes; AG, Aargau; BE, Bern; JU, Jura; LU, Luzern; SG, St Gallen; TG, Thurgau; ZH, Zurich). and numbers indicate individual farms.



to strain (Table 2). The ST1-t2279-IVc isolate was only detected once and showed resistance to  $\beta$ -lactams [*mecA*, *blaZ*], tetracycline [*tet(M)*], MLS<sub>B</sub> antibiotics [*erm(A)*], and spectinomycin [*ant(9)-Ia*]; additionally, a resistance to ciprofloxacin was detected (Table 2). The

resistance mechanism to fluoroquinolones which may be due to mutations in the topoisomerase genes [27] was not determined. In contrast, MRSA isolates from 2009 and 2010 were susceptible to vancomycin, gentamicin, kanamycin, chloramphenicol, quinupristin/dalfopristin,

**Table 2 The clonal lineage and resistance profile of methicillin-resistant *Staphylococcus aureus* from pig noses in Switzerland**

MLST	spa type	SCCmec type	Number of isolates	Farm <sup>a</sup>	Representative strain	Antibiotic resistance genes and resistance breakpoints (mg/L) <sup>b</sup>										
						FOX	PEN	TET	ERY	CLI	STR	TIA	SPC	TMP	SMX	CIP
						(>4)	(>0.125)	(>2)	(>2)	(>0.5)	(>16)	(>2)	(>128)	(>2)	(>128)	(>1)
<b>2009</b>																
ST1	t2279	IVc	1	TG3	IMD887-09	<i>mecA</i>	<i>mecA</i> <i>blaZ</i>	<i>tet(M)</i>	<i>erm</i> (A)	<i>erm</i> (A)			<i>ant(9)-la</i>			+
ST398	t011	V	1	SG1	IMD358-09	<i>mecA</i>	<i>mecA</i>	<i>tet(M)</i> <i>tet(K)</i>				<i>vga</i> (A)v				
ST398	t011	V	1	TG1	IMD355-09	<i>mecA</i>	<i>mecA</i>	<i>tet(M)</i> <i>tet(K)</i>	<i>erm</i> (C)	<i>erm</i> (C)			+			
ST398	t011	V	1	LU2	IMD1753-09	<i>mecA</i>	<i>mecA</i>	<i>tet(M)</i> <i>tet(K)</i>					+			
ST398	t034	V	1	BE1	IMD1116-09	<i>mecA</i>	<i>mecA</i> <i>blaZ</i>	<i>tet(M)</i> <i>tet(K)</i>	<i>erm</i> (A)	<i>erm</i> (A)	<i>str</i>	+	<i>ant(9)-la</i>	<i>dfr</i> (G)		
ST398	t1451	V	1	LU1	IMD1270-09	<i>mecA</i>	<i>mecA</i>	<i>tet(M)</i> <i>tet(K)</i>	<i>erm</i> (C)	<i>erm</i> (C)	<i>str</i>	<i>vga</i> (A)v				
ST49	t208	V	1	TG2	IMD1000-09	<i>mecA</i>	<i>mecA</i>	<i>tet(M)</i> <i>tet(K)</i>	<i>erm</i> (C)	<i>erm</i> (C)		<i>vga</i> (A)v				+
ST49	t208	V	1	SG2	IMD1771-09	<i>mecA</i>	<i>mecA</i>	<i>tet(M)</i> <i>tet(K)</i>				<i>vga</i> (A)v				+
<b>2010</b>																
ST398	t034	V	16	TG5 TG8 TG9 TG10 LU3 LU4 LU5 LU6 BE2 BE3 BE4 AG1 JU1 SG3 SG5 ZH1	IMD49-10	<i>mecA</i>	<i>mecA</i> <i>blaZ</i>	<i>tet(M)</i> <i>tet(K)</i>	<i>erm</i> (A)	<i>erm</i> (A)	<i>str</i>	+	<i>ant(9)-la</i>	<i>dfr</i> (G)		
ST398	t034	V	1	TG 7	IMD704-10	<i>mecA</i>	<i>mecA</i> <i>blaZ</i>	<i>tet(M)</i> <i>tet(K)</i>	<i>erm</i> (A)	<i>erm</i> (A)		+	<i>ant(9)-la</i>	<i>dfr</i> (G)		
ST398	t011	V	1	AI1	IMD233-10	<i>mecA</i>	<i>mecA</i>	<i>tet(M)</i> <i>tet(K)</i>	<i>erm</i> (C)	<i>erm</i> (C)			+			
ST49	t208	V	3	SG4 SG6 TG4	IMD426-10	<i>mecA</i>	<i>mecA</i>	<i>tet(M)</i> <i>tet(K)</i>				<i>vga</i> (A)v				+
ST49	t208	V	1	TG6	IMD603-10	<i>mecA</i>	<i>mecA</i>	<i>tet(M)</i> <i>tet(K)</i>	<i>erm</i> (A)	<i>erm</i> (A)		<i>vga</i> (A)v				+
ST49	t208	V	1	AG2	IMD2002-10	<i>mecA</i>	<i>mecA</i>	<i>tet(M)</i> <i>tet(K)</i>			<i>str</i>	<i>vga</i> (A)v				+

a) Geographical origin of the farm: letters indicate the Cantons (AI, Appenzell Innen Rhodes; AG, Aargau; BE, Bern; JU, Jura; LU, Luzern; SG, St Gallen; TG, Thurgau; ZH, Zurich), and numbers indicate individual farms.

b) CIP, ciprofloxacin; CLI, clindamycin; ERY, erythromycin; FOX, cefoxitin; PEN, penicillin; SPC, spectinomycin; SMX, sulphamethoxazole; STR, streptomycin; TET, tetracycline; TIA, tiamulin; TMP, trimethoprim. The MIC breakpoints (in micrograms per millilitre) that determine resistance were recommended from EUCAST for *S. aureus* <http://www.eucast.org>. Resistance breakpoints for tiamulin, spectinomycin, streptomycin and sulphamethoxazole were tentatively derived from epidemiological MIC cut-off values from EUCAST.

Antibiotic resistance genes and their functions are indicated as follows: *ant(9)-la*, spectinomycin adenylnucleotidyltransferase; *blaZ*, β-lactamase; *dfr(G)*, dihydrofolate reductase; *erm(A)* and *erm(C)*, macrolides, lincosamides and streptogramins B rRNA methylase; *mecA*, penicillin-binding protein PBP2a; *str*, streptomycin adenylnucleotidyltransferase; *tet(K)*, tetracycline efflux protein; *tet(M)*, tetracycline ribosomal protection protein; *vga (A)v*, pleuromutilins and streptogramins A ATP binding transporter.

ND, not detected; +, The MIC values were greater than the resistance breakpoint, but the resistance mechanism remained uncharacterised; blank spaces indicate no resistance



fusidic acid, mupirocin, rifampicin, and linezolid; importantly, none of the isolates carried the Panton-Valentine Leukocidin toxin.

## Discussion

### Molecular typing and epidemiology

This study was the first to reveal the presence of MRSA ST49 in pigs. MRSA ST49 was already present in 2009, but its prevalence increased in 2010. The presence of the new clonal lineage ST49-t208, which is unique to MRSA from pigs in Switzerland, suggests that the selection occurred within the Swiss pig population. This idea is supported by the fact that MRSA ST49 was detected in seven different pigs from seven different farms slaughtered in three different slaughterhouses, and that ST49 isolates displayed different resistance profiles mainly due to the acquisition of an additional *erm(A)* or *erm(C)* gene (Table 2). The *spa* type t208, which is associated with ST49, was previously found in seven out of 48 methicillin-susceptible *S. aureus* MSSA from pigs in Switzerland [20], suggesting a possible acquisition of the *SCCmec* element. ST49 is rarely mentioned in the literature; it was not detected among 2890 MSSA and MRSA from human infections in the 26 EU countries, the 133 *S. aureus* colonising healthy adults in Switzerland nor within the 572 MSSA isolates from Swiss children [28-30]. ST49 has only been associated with MSSA involved in one human infection in the United Kingdom and in three cases of skin lesions and laryngeal ulceration in wild squirrels [31,32]. In the latter three cases, the authors mentioned close contact with persons who fed the squirrels as a possible source of infection. Further screening of humans, including people working with food-producing animals, would be necessary to determine if humans also play a role in the dissemination of MRSA ST49, as demonstrated with ST398. Human carriage of ST398 has been widely documented in other European countries, and the exchange of ST398 between humans and pigs has been reported in the Netherlands and Denmark [13,33,34]. Furthermore, ST398-t034 was the most predominant lineage found in Swiss pigs, and has been detected in one out of 133 veterinarians in Switzerland [22]. In our study, ST1 was only detected once, although it has been associated with MRSA infections in humans in the United Kingdom and Spain, and in horses in Austria [31,35,36]. ST1 represented, however, 15.6% of the MRSA-positive pig finishing holdings in Italy [37]. Cases of transmission of ST1 between humans and cows have also been reported [38].

Transmission of MRSA among pigs during animal transport and at slaughterhouse has been described [39]. In our study, no direct association of STs to a specific abattoir has been observed, except for slaughterhouse A, where ST398-t034 has been predominantly detected in

the samples. Transmission which may have occurred at slaughterhouse may not be excluded for this abattoir. Otherwise, MRSA ST398 and ST49 were found in different slaughterhouses at different sampling dates and the pigs originated from different farms.

### Increase of MRSA prevalence

This study is also the first to describe MRSA prevalence in slaughter pigs in Switzerland using a sampling plan which was representative of the Swiss slaughter pig population and which was evenly distributed over time. This strategy allowed us to obtain statistically significant and representative data for the evaluation of the chronological distribution of MRSA. Such representative sampling criteria were not considered in the two previous studies reporting MRSA in pigs in Switzerland, resulting in a less accurate estimation of the prevalence over the entire country [20,22]. However, these studies indicated that the prevalence of MRSA was low among the proportion of pigs analysed before 2009. Our study also showed that the prevalence of MRSA in 2009 was low, and indicated an approximate three-fold increase within one year. The increased diversity of MRSA and the emergence of the clonal lineage ST49 unique to MRSA from Swiss pigs suggested that MRSA-positive pigs imported into Switzerland have not introduced this MRSA. However, Switzerland sporadically imports a low number of breeding pigs, from which MRSA ST398 may have been introduced. In 2010, for example, Switzerland imported pigs from Denmark ( $n = 10$ ), The Netherlands ( $n = 4$ ), and Germany ( $n = 1$ ), where the prevalence of MRSA ST398 is high (Federal Veterinary Office, pers. communication). Like Switzerland, the predominant clone in Denmark and Germany is ST398-t034, whereas MRSA ST398-t011 is the predominant in The Netherlands [40-42].

## Conclusions

Two predominant clonal lineages, ST398-t034 and ST49-t208, are spreading within the Swiss slaughter pig population. Focus should be stressed on prevention of the transmission of these multi-resistant zoonotic bacteria from the farms into the community. Human carriage of MRSA represents a new challenge for public health because people that are in the vicinity of animals have been shown to have a higher risk of developing a MRSA infection when hospitalised. Targeted screening for MRSA in at-risk people, that is, farmers, veterinarians, and other people with close contact to animals, as recommended by Harbarth et al. (2011) [43], should be taken into account upon hospital admission. A study on MRSA decolonisation in humans with close contact to pigs showed a low efficacy due to continuous re-colonisation [44]. Therefore, effective hygiene measures into the entire pig production chain should be maintained as proposed by the Scientific

Advisory Group on Antimicrobials [8]. Nevertheless, periodic monitoring of the MRSA in livestock animals at slaughterhouses is now established in Switzerland at the National Reference Laboratory for Antimicrobial Resistance (Centre of Zoonoses, Bacterial Animal Diseases and Antimicrobial Resistance (ZOBA), Institute of Veterinary Bacteriology, Vetsuisse Faculty, University of Bern), for the surveillance of MRSA in the livestock population.

## Methods

### Sampling

Representative samples were taken in accordance with the framework of a national monitoring program on antimicrobial resistance in food animals [23]. A minimum required sample size of 382 randomly selected animals was calculated with the assumptions of an infinite population size, a prevalence of 5%, a desired confidence level of 97.5% and an accuracy of 2.5%. The samples were randomly taken at the nine biggest abattoirs (A to I), where over 85% of the pigs in Switzerland were slaughtered. Only one sample was taken per animal holding. The number of samples in the sampling frame collected from each slaughterhouse was proportional to the number of pigs slaughtered at each establishment per year. One to 8 samples were taken per sampling day and abattoir. The samples collected on Mondays and Tuesdays were equally distributed over the years. Based on these data, random sampling plans were conducted, resulting in a total of 405 nasal swabs of fattening pigs in 2009 and 392 swabs from pigs in 2010.

### Isolation of MRSA

Samples were taken using transport swabs (Oxoid Ltd, Basingstoke, England) from the nares of the pigs subsequent to stunning by officials of the Swiss abattoir authorities and immediately after sampling, were transported to the laboratory without cooling. After arrival, swabs were transferred into tubes containing 10 ml Mueller Hinton Broth supplemented with 6.5% NaCl and incubated aerobically at 37°C for 24 h under agitation. One ml from this pre-enrichment was inoculated into 9 ml tryptone soy broth containing 3.5 mg/L cefoxitin and 75 mg/L aztreonam, and further incubated aerobically at 37°C for 24 h. A loopful was then spread onto MRSA selective agar plates (BBL™ CHROMagar™ MRSA; Becton Dickinson, Franklin Lakes, NJ), which were incubated at 37°C for 24 h. Pink to mauve-colored colonies were regarded as suspicious and five presumptive colonies were cultivated onto tryptone soy agar plates containing 5% sheep blood (TSA-SB) (Oxoid Ltd, Basingstoke, England) at 37°C for 24 h. *S. aureus* was identified using Vitek 2 with Gram-Positive (GP) cards (BioMérieux, Mary l'Etoile, France) following manufacturer's recommendations.

### Molecular typing and antibiotic resistance

The minimal inhibitory concentration (MIC) of the antibiotics was determined by broth microdilution in Müller-Hinton using the Sensititre susceptibility plate EUST (Trek Diagnostics Systems, East Grinstead, England; MCS Diagnostics BV, Swalmen, The Netherlands), except for spectinomycin, which was tested using homemade microbroth dilution plates. The MIC was interpreted according to the European Committee on Antimicrobial Susceptibility Testing (EUCAST) guidelines <http://www.eucast.org> using clinical resistance breakpoints for cefoxitin (> 4 mg/L), chloramphenicol (> 8 mg/L), clindamycin (> 0.5 mg/L), ciprofloxacin (> 1 mg/L), erythromycin (> 2 mg/L), fusidic acid (> 1 mg/L), gentamicin (> 1 mg/L), linezolid (> 4 mg/L), penicillin (> 0.125 mg/L), quinupristin/dalfopristin (> 2 mg/L), rifampicin (> 0.5 mg/L), tetracycline (> 2 mg/L), trimethoprim (> 4 mg/L), and vancomycin (> 2 mg/L). Otherwise, resistance breakpoints were tentatively derived from epidemiological MIC cut off values from EUCAST for kanamycin (> 8 mg/L), mupirocin (> 1 mg/L), streptomycin (> 16 mg/L), spectinomycin (> 128 mg/L), sulfamethoxazole (> 128 mg/L), and tiamulin (> 2 mg/L). Antibiotic resistance genes, including *mecA*, were detected using a microarray [45]. SCC*mec* types were determined by multiplex Polymerase Chain Reaction PCR assays [46]. Sequence types (ST) were determined using multilocus sequence typing (MLST) [31]. *Spa* type was determined as previously described and analysed using the Ridom StaphType software (Ridom StaphType, Ridom GmbH, Würzburg, Germany) [47]. The presence of the Pantone-Valentine Leukocidin operon *lukS-lukF* was determined by PCR as described previously [48].

### Statistical Analysis

Sampling calculations were performed using the Win Episcope 2.0 software <http://www.clive.ed.ac.uk/winepiscope>. The difference in the prevalence between the years was analysed by Chi-square test. The level of significance was 5%. All statistics were performed using the NCSS 2007 statistical software (Hintze J. NCSS (2008) Version 07.1.8. Kaysville, UT, USA).

### Acknowledgements

We thank the participating slaughterhouses and their personnel for the sampling and acquisition of data. We also thank E. Lüthi, I. Brodard, and M. Binggeli for their technical assistance. The Swiss Federal Veterinary Office and the Canton of Bern financed this study.

### Author details

<sup>1</sup>Institute of Veterinary Bacteriology, Vetsuisse Faculty, University of Bern, Länggass-Strasse 122, CH-3012 Bern, Switzerland. <sup>2</sup>Centre of Zoonoses, Animal Bacterial Diseases and Antibiotic Resistance (ZOBA), Institute of Veterinary Bacteriology, Vetsuisse Faculty, University of Bern, Länggass-Strasse 122, CH-3012 Bern, Switzerland. <sup>3</sup>Federal Veterinary Office, Schwarzenburgstrasse 155, CH-3003 Bern, Switzerland.

#### Authors' contributions

GO wrote the manuscript, supervised the analyses for the detection and identification of MRSA and contributed to the conception of the study. SB designed the sampling plan, organised the sampling at the slaughterhouses and performed statistical analyses. AR carried out the molecular genetic studies. VP designed and coordinated the study, and helped to write the manuscript. All authors read and approved the final manuscript.

Received: 9 March 2011 Accepted: 24 June 2011

Published: 24 June 2011

#### References

- Cuny C, Friedrich A, Kozytska S, Layer F, Nubel U, Ohlsen K, et al: **Emergence of methicillin-resistant *Staphylococcus aureus* (MRSA) in different animal species.** *Int J Med Microbiol* 2010, **300**:109-117.
- Weese JS: **Methicillin-resistant *Staphylococcus aureus* in animals.** *ILAR J* 2010, **51**:233-244.
- Vanderhaeghen W, Hermans K, Haesebrouck F, Butaye P: **Methicillin-resistant *Staphylococcus aureus* (MRSA) in food production animals.** *Epidemiol Infect* 2010, **138**:606-625.
- Smith TC, Pearson N: **The emergence of *Staphylococcus aureus* ST398.** *Vector Borne Zoonotic Dis* 2011, **11**:327-339.
- Wagenaar JA, Yue H, Pritchard J, Broekhuizen-Stins M, Huijsdens X, Mevius DJ, et al: **Unexpected sequence types in livestock associated methicillin-resistant *Staphylococcus aureus* (MRSA): MRSA ST9 and a single locus variant of ST9 in pig farming in China.** *Vet Microbiol* 2009, **139**:405-409.
- Guardabassi L, O'Donoghue M, Moodley A, Ho J, Boost M: **Novel lineage of methicillin-resistant *Staphylococcus aureus*, Hong Kong.** *Emerg Infect Dis* 2009, **15**:1998-2000.
- Vanderhaeghen W, Clerpentier T, Adriaensen C, Vicca J, Hermans K, Butaye P: **Methicillin-resistant *Staphylococcus aureus* (MRSA) ST398 associated with clinical and subclinical mastitis in Belgian cows.** *Vet Microbiol* 2010, **144**:166-171.
- Catry B, van Duijkeren E, Pomba MC, Greko C, Moreno MA, Pyorala S, et al: **Reflection paper on MRSA in food-producing and companion animals: epidemiology and control options for human and animal health.** *Epidemiol Infect* 2010, **138**:626-644.
- Wulf M, Voss A: **MRSA in livestock animals-an epidemic waiting to happen?** *Clin Microbiol Infect* 2008, **14**:519-521.
- van Loo I, Huijsdens X, Tiemersma E, de Neeling A, van de Sande-Bruinsma N, Beaujean D, et al: **Emergence of methicillin-resistant *Staphylococcus aureus* of animal origin in humans.** *Emerg Infect Dis* 2007, **13**:1834-1839.
- Mulders MN, Haenen AP, Geenen PL, Vesseur PC, Poldervaart ES, Bosch T, et al: **Prevalence of livestock-associated MRSA in broiler flocks and risk factors for slaughterhouse personnel in The Netherlands.** *Epidemiol Infect* 2010, **138**:743-755.
- Khanna T, Friendship R, Dewey C, Weese JS: **Methicillin resistant *Staphylococcus aureus* colonization in pigs and pig farmers.** *Vet Microbiol* 2008, **128**:298-303.
- van Belkum A, Melles DC, Peeters JK, van Leeuwen WB, van Duijkeren E, Huijsdens XW, et al: **Methicillin-resistant and-susceptible *Staphylococcus aureus* sequence type 398 in pigs and humans.** *Emerg Infect Dis* 2008, **14**:479-483.
- Smith TC, Male MJ, Harper AL, Kroeger JS, Tinkler GP, Moritz ED, et al: **Methicillin-resistant *Staphylococcus aureus* (MRSA) strain ST398 is present in midwestern U.S. swine and swine workers.** *PLoS ONE* 2009, **4**: e4258.
- van Rijen MML, van Keulen PH, Kluytmans JA: **Increase in a Dutch Hospital of methicillin-resistant *Staphylococcus aureus* related to animal farming.** *Clin Infect Dis* 2008, **46**:261-263.
- van Rijen MM, Bosch T, Heck ME, Kluytmans JA: **Methicillin-resistant *Staphylococcus aureus* epidemiology and transmission in a Dutch hospital.** *J Hosp Infect* 2009, **72**:299-306.
- Witte W, Strommenger B, Stanek C, Cuny C: **Methicillin-resistant *Staphylococcus aureus* ST398 in humans and animals, Central Europe.** *Emerg Infect Dis* 2007, **13**:255-258.
- Potel C, Alvarez-Fernandez M, Constenla L, Alvarez P, Perez S: **First human isolates of methicillin-resistant *Staphylococcus aureus* sequence type 398 in Spain.** *Eur J Clin Microbiol Infect Dis* 2010, **29**:351-352.
- European Food Safety Authority: **Analysis of the baseline survey on the prevalence of methicillin resistant *Staphylococcus aureus* (MRSA) in holding with breeding pigs, in the EU, 2008, Part A: MRSA prevalence estimates.** *EFSA J* 2009, **7**:1376.
- Riesen A, Perreten V: **Antibiotic resistance and genetic diversity in *Staphylococcus aureus* from slaughter pigs in Switzerland.** *Schweiz Arch Tierheilk* 2009, **151**:425-431.
- Nitzsche S, Zweifel C, Stephan R: **Phenotypic and genotypic traits of *Staphylococcus aureus* strains isolated from pig carcasses.** *Vet Microbiol* 2007, **120**:292-299.
- Huber H, Koller S, Giezendanner N, Stephan R, Zweifel C: **Prevalence and characteristics of methicillin-resistant *Staphylococcus aureus* in humans in contact with farm animals, in livestock, and in food of animal origin, Switzerland, 2009.** *Euro Surveill* 2010, **15**:1-4.
- Büttner S, Kuhn M: **Antibiotikaresistenzmonitoring-Jahresbericht. 2008** [http://www.bvet.admin.ch/themen].
- Jung YH, Shin ES, Kim O, Yoo JS, Lee KM, Yoo JI, et al: **Characterization of two newly identified genes, *vgaD* and *vatG*, conferring resistance to streptogramin A in *Enterococcus faecium*.** *Antimicrob Agents Chemother* 2010, **54**:4744-4749.
- Long KS, Poehlsgaard J, Kehrenberg C, Schwarz S, Vester B: **The Cfr rRNA methyltransferase confers resistance to Phenicolis, Lincosamides, Oxazolidinones, Pleuromutilins, and Streptogramin A antibiotics.** *Antimicrob Agents Chemother* 2006, **50**:2500-2505.
- Miller K, Dunsmore CJ, Fishwick CW, Chopra I: **Linezolid and tiamulin cross-resistance in *Staphylococcus aureus* mediated by point mutations in the peptidyl transferase center.** *Antimicrob Agents Chemother* 2008, **52**:1737-1742.
- Hooper DC: **Mechanisms of action and resistance of older and newer fluoroquinolones.** *Clin Infect Dis* 2000, **31**(Suppl 2):S24-S28.
- Grundmann H, Aanensen DM, van den Wijngaard CC, Spratt BG, Harmsen D, Friedrich AW: **Geographic distribution of *Staphylococcus aureus* causing invasive infections in Europe: a molecular-epidemiological analysis.** *PLoS Med* 2010, **7**:e1000215.
- Sakwinska O, Kuhn G, Balmelli C, Francioli P, Giddey M, Perreten V, et al: **Genetic diversity and ecological success of *Staphylococcus aureus* strains colonizing humans.** *Appl Environ Microbiol* 2009, **75**:175-183.
- Megevang C, Gervaix A, Heiningen U, Berger C, Aebi C, Vaudaux B, et al: **Molecular epidemiology of the nasal colonization by methicillin-susceptible *Staphylococcus aureus* in Swiss children.** *Clin Microbiol Infect* 2010, **16**:1414-1420.
- Enright MC, Day NP, Davies CE, Peacock SJ, Spratt BG: **Multilocus sequence typing for characterization of methicillin-resistant and methicillin-susceptible clones of *Staphylococcus aureus*.** *J Clin Microbiol* 2000, **38**:1008-1015.
- Simpson V, Davison N, Hudson L, Whatmore AM: ***Staphylococcus aureus* ST49 infection in red squirrels.** *Vet Rec* 2010, **167**:69.
- Otter JA, French GL: **Molecular epidemiology of community-associated methicillin-resistant *Staphylococcus aureus* in Europe.** *Lancet Infect Dis* 2010, **10**:227-239.
- Lewis HC, Mølbak K, Reese C, Aarestrup FM, Selchau M, Sørup M, et al: **Pigs as source of methicillin-resistant *Staphylococcus aureus* CC398 infections in humans, Denmark.** *Emerg Infect Dis* 2008, **14**:1383-1389.
- Aspiroz C, Lozano C, Vindel A, Lasarte JJ, Zarazaga M, Torres C: **Skin lesion caused by ST398 and ST1 MRSA, Spain.** *Emerg Infect Dis* 2010, **16**:157-159.
- Cuny C, Strommenger B, Witte W, Stanek C: **Clusters of infections in horses with MRSA ST1, ST254, and ST398 in a veterinary hospital.** *Microb Drug Resist* 2008, **14**:307-310.
- Battisti A, Franco A, Meriardi G, Hasman H, Iurescia M, Lorenzetti R, et al: **Heterogeneity among methicillin-resistant *Staphylococcus aureus* from Italian pig finishing holdings.** *Vet Microbiol* 2010, **142**:361-366.
- Juhász-Kaszanyitzky E, Janosi S, Somogyi P, Dan A, van der Graaf-van Bloois, van Duijkeren E, et al: **MRSA transmission between cows and humans.** *Emerg Infect Dis* 2007, **13**:630-632.
- Broens EM, Graat EAM, van der Wolf PJ, van de Giessen AW, De Jong MCM: **Transmission of methicillin resistant *Staphylococcus aureus* among pigs during transportation from farm to abattoir.** *Vet J* 2010.
- de Neeling AJ, van den Broek MJM, Spalburg EC, van Santen-Verheuevel MG, Dam-Deisz WDC, Boshuizen HC, et al: **High prevalence of methicillin resistant *Staphylococcus aureus* in pigs.** *Vet Microbiol* 2007, **122**:366-372.



41. Hasman H, Moodley A, Guardabassi L, Stegger M, Skov RL, Aarestrup FM: **Spa type distribution in *Staphylococcus aureus* originating from pigs, cattle and poultry.** *Vet Microbiol* 2010, **141**:326-331.
42. Tenhagen BA, Fetsch A, Stuhrenberg B, Schleuter G, Guerra B, Hammerl JA, *et al*: **Prevalence of MRSA types in slaughter pigs in different German abattoirs.** *Vet Rec* 2009, **165**:589-593.
43. Harbarth S, Hawkey PM, Tenover F, Stefani S, Pantosti A, Struelens MJ: **Update on screening and clinical diagnosis of methicillin-resistant *Staphylococcus aureus* (MRSA).** *Int J Antimicrob Agents* 2011, **37**:110-117.
44. Lozano C, Aspiroz C, Lasarte JJ, Gomez-Sanz E, Zarazaga M, Torres C: **Dynamic of nasal colonization by methicillin-resistant *Staphylococcus aureus* ST398 and ST1 after mupirocin treatment in a family in close contact with pigs.** *Comp Immunol Microbiol Infect Dis* 2011, **34**:e1-e7.
45. Perreten V, Vorlet-Fawer L, Slickers P, Ehricht R, Kuhnert P, Frey J: **Microarray-based detection of 90 antibiotic resistance genes of gram-positive bacteria.** *J Clin Microbiol* 2005, **43**:2291-2302.
46. Kondo Y, Ito T, Ma XX, Watanabe S, Kreiswirth BN, Etienne J, *et al*: **Combination of multiplex PCRs for Staphylococcal Cassette Chromosome *mec* type assignment: rapid identification system for *mec*, *ccr*, and major differences in junkyard regions.** *Antimicrob Agents Chemother* 2007, **51**:264-274.
47. Harmsen D, Claus H, Witte W, Rothgänger J, Claus H, Turnwald D, *et al*: **Typing of methicillin-resistant *Staphylococcus aureus* in a university hospital setting by using novel software for *spa* repeat determination and database management.** *J Clin Microbiol* 2003, **41**:5442-5448.
48. Lina G, Piémont Y, Godail-Gamot F, Bes M, Peter M-O, Gauduchon V, *et al*: **Involvement of Panton-Valentine leukocidin-producing *Staphylococcus aureus* in primary skin infections and pneumonia.** *Clin Infect Dis* 1999, **29**:1128-1132.

doi:10.1186/1746-6148-7-30

**Cite this article as:** Overesch *et al.*: The increase of methicillin-resistant *Staphylococcus aureus* (MRSA) and the presence of an unusual sequence type ST49 in slaughter pigs in Switzerland. *BMC Veterinary Research* 2011 **7**:30.

**Submit your next manuscript to BioMed Central  
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)

