

RESEARCH

Open Access



Phenotypic, molecular detection and antibiogram analysis of *Aeromonas Hydrophila* from *Oreochromis Niloticus* (Nile Tilapia) and Ready-To-eat fish products in selected Rift Valley lakes of Ethiopia

Nebiyu Kassa Kerigano^{1*}, Tesfaye Rufael Chibsa², Yitbarek Getachew Molla³, Abde Aliy Mohammed⁴, Mekdes Tamiru⁵, Abebe Olani Bulto⁵, Tafesse Koran Wodaj⁵, Dereje Shegu Gebreweld⁵ and Alemu Kebede Abdi^{1*}

Abstract

Background *Aeromonas hydrophila* is a zoonotic bacterial pathogen that frequently causes disease and mass mortalities among cultured and feral fishes worldwide. In Ethiopia, *A. hydrophila* outbreak was reported in Sebeta fish ponds and in Lake Tana fishery. However, there is no to little information on the molecular, and phenotypical characteristics of *A. hydrophila* in Ethiopian fisheries. Therefore, a cross-sectional study was conducted from November 2020 to May 2021 in selected Ethiopian Rift valley lakes.

Results A total of 140 samples were collected aseptically from fish (Muscle, Gill, Intestine, Spleen and Kidney) from fish landing sites, market and restaurants with purposive sampling methods. *Aeromonas* selective media (AMB), morphological and biochemical tests were used to isolate and identify *A. hydrophila*. Accordingly, the pathogen was isolated from 81 (60.45%) of samples. Among the isolates 92.59% expressed virulence trait through β hemolysis on blood agar media with 5% sheep blood. Moreover, 54 strains (66.67%) were further confirmed with Real-Time PCR (qPCR) using ahal gene specific primers and optimized protocol. The highest (68.51%) were detected from live fish, (24.07%) were from market fish and the lowest (7.4%) were from ready-to-eat products. Antibiogram analysis was conducted on ten representative isolates. Accordingly, *A. hydrophila* isolates were susceptible to ciprofloxacin (100%), chloramphenicol (100%) and ceftriaxone (100%). However, all ten isolates were resistant to Amoxicillin and Penicillin.

Conclusions The study indicates *A. hydrophila* strains carrying virulence ahal gene that were β -hemolytic and resistant to antibiotics commonly used in human and veterinary medicine are circulating in the fishery. The detection of the pathogen in 140 of the sampled fish population is alarming for potential outbreaks and zoonosis. Therefore, further molecular epidemiology of the disease should be studied to establish potential inter host transmission and

*Correspondence:

Nebiyu Kassa Kerigano
neby.etu@gmail.com; k.nebiyu@ahi.gov.et

Full list of author information is available at the end of the article



© The Author(s) 2023. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

antibiotic resistance traits. Therefore, raising the public awareness on risk associated with consuming undercooked or raw fish meat is pertinent.

Keywords *A. hydrophila*, Ethiopia, Nile tilapia, qPCR, Rift valley lake

Background

Nile tilapia (*Oreochromis niloticus*) is one of the commercially important fast-growing and well adapted freshwater fish that is produced extensively and intensively all over the world [1]. Tilapia are increasingly used in aquaculture and is currently the second most important freshwater fish farmed worldwide with an annual global production of 6.4 MT [2]. It is characterized by their reasonable resistance to diseases and its suitability for intensive farming which subsequently leads to increased production and makes it as a cheap protein source for all people [3]. Nowadays this high protein source is threatened by bacterial diseases especially those caused by drug resistance and highly virulent bacteria such as *A. hydrophila* [4]. Significantly impeding both economic and socioeconomic developments in regions dependent on aquaculture and fisheries and zoonotic implications as well [5].

A. hydrophila is facultative anaerobic, Gram-negative bacteria that belong to the family Aeromonadaceae which is cosmopolitan in distribution and have a broad host spectrum with both cold and warm blooded animals including humans [6]. *A. hydrophila* is a wellknown bacterial pathogen that frequently causes disease and mass mortalities among cultured and feral fishes worldwide [7]. *A. hydrophila* has gained increased attention due to pathogenicity to humans and emerged as a foodborne pathogen of extreme importance [8]. *A. hydrophila* resulting serious health condition and death associated with consumption of frozen fish in market-sold sushi products containing raw fish [9]. High antibiotic resistance is seen in *A. hydrophila* infections [10] and regarded universally exhibit resistance to the penicillin for quite a long time [11] nowadays, becoming a serious public health concern. In Ethiopia however, less attention has been given to pathogens of fish including those which have zoonotic importance except few isolated cases [12]. For instance, a survey of bacterial and parasitic fish pathogens was conducted in Lake Ziway but *A. hydrophila* was not included [13]. *A. hydrophila* was reported as the most frequent isolate from Lake Tana and also the pathogen was associated with outbreak and mortality in Sebeta fish ponds [14].

In Ethiopia, intensive and semi-intensive aquaculture is becoming an emerging business in the country. The number of private investors interested in fish farming in the country is evolving and some of them have even already started the process. The Great Renaissance Dam and several other dams and reservoirs are being constructed

in the country for hydropower generation, irrigation and other purposes apart from providing water for their primary uses, these water bodies could also be stocked with different fish species which could provide a source of livelihood to many rural young Ethiopians engaged in fishing. Despite the potential contribution of fisheries in the country emerging zoonotic bacterial pathogen like *A. hydrophila* could constrain the productivity and safety of the fish industry in the country. This calls for proactive investigation into important pathogens in water bodies with high fish sources in Rift Valley lakes of Ethiopia.

According to FAO [1], majority of fish catch in Ethiopia originate from Rift valley lakes. Therefore, knowing the infection status and characteristics of *A. hydrophila* in fish and ready-to-eat fish products is paramount to the understanding of the epidemiology and associated risks to public health. To this end the present study was intended to isolate and determine phenotypic and genotypic features of *A. hydrophila* infecting tilapia in selected Rift Valley Lakes and fish products in respective towns. The specific objectives of the study were to isolate *A. hydrophila* from fish and ready-to-eat fish products, to determine the susceptibility of *A. hydrophila* isolates to major antimicrobials of veterinary and human importance and to reveal phenotypic and genotypic traits of *A. hydrophila* isolates.

Results

Clinical and post-mortem findings

Fishes suspected of infection with *A. hydrophila* showed hemorrhages all over the body especially at the base of fins and tail. Clinical presentations observed include fins rot, cloudiness of both eyes, detachment of scales and skin ulceration and abdominal distention. Internally these fishes showed abdominal dropsy with reddish ascetic exudates, liver paleness and enlargement in some fishes and congested with necrotic patches in other fishes, spleen was congested, enlarged and hemorrhagic enteritis in some fishes as shown in (Fig. 1).

An arrow in (A) shows abdominal dropsy with reddish ascetic exudates (B) shows skin hemorrhage at the base of pectoral fin with hemorrhagic skin ulcer under the dorsal and tail fin and dark discoloration in the skin (C) shows skin ulcer.

Bacteriological identification and biochemical characterization of *A. hydrophila*

The presumptive identification of the bacteria in the current study was carried out from the colony morphology




Fig. 1 Clinical picture and post mortem findings




Fig. 2 Characters of *A. hydrophila* on (A) Aeromonas medium base (B) Mac-Concey agar (C) Nutrient Agar

over Aeromonas Medium Base, a selective medium for *A. hydrophila*. Accordingly, based on 14 morphological and biochemical tests, a total number of 81(60.45%) isolates were presumptively identified as *A. hydrophila*. They appeared rounded smooth colonies 2-3 mm in diameter and dark green with a darker center in Aeromonas medium base, pale like shaped on MacConkey agar indicated that *A. hydrophila* is unable to ferment lactose sugar and creamy white on Nutrient agar as presented in (Fig. 2). Colonies were gram-negative short rods, they gave a positive reaction for oxidase, catalase, DNase, Indole production, also ferment glucose with production of acid and gas, sugar utilization K/A, Acid production from (Sucrose and Mannitol) and Motile. They gave negative results toward xylose, urea hydrolysis, and non-lactose fermentation and produced variable results with MRVP.

Hemolysis assay

Hemolytic activity of the isolates was determined for its importance as a virulent factor. *A. hydrophila* produced hemolysis on blood agar base with 5% sheep blood. Accordingly, from the current study found that 93.33% (n=56/60), 94.11% (n=16/17) and 75% (n=3/4) isolates from the life fish group, market fish and RTE fish show β hemolysis respectively and 6.66% (n=4/60), 5.88% (n=1/17) and 25% (n=1/4) show α hemolysis. The hemolysis pattern results in the media displaying clear halos around bacterial colonies as shown in (Fig. 3). Hemolytic activities of *A. hydrophila* from the current study found over all isolates 92.59% (n=75/81) show β hemolysis and only 7.4% (n=6/81) of α hemolysis as shown in (Table 1).




Fig. 3 β -hemolysis activity by *A. hydrophila* on blood agar base with 5% sheep blood

Table 1 Hemolytic characteristics of the isolates

Hemolytic activity of <i>A. hydrophila</i> isolated from fish samples			
Source	Total	β	α
Live fish	60	56 (93.33)	4 (6.66)
Market fish	17	16 (94.11)	1 (5.88%)
RTE	4	3 (75%)	1 (25%)
Total	81	75 (92.59%)	6 (7.4%)

β : beta, α : alpha

Molecular detection

Quantitative real-time PCR detection of *A. hydrophila* and virulence gene

Molecular detection with Real-Time PCR (qPCR) using specific primers based on the sequence of the *ahaI* gene coding for adhesive surface protein mainly present in virulent *A. hydrophila* strain. From the total of 81 *A. hydrophila* isolates, 54 were confirmed by real-time PCR for presence of the *ahaI* gene. The threshold cut off value for classifications of the samples as positive or negative by the real-time PCR was set to a cycle threshold (Ct) value of 34. Samples giving a Ct value of ≤ 34 with a sigmoid shape of the analysis curve were classified as positive (Fig. 4). Samples with a Ct value > 34 were classified as negative. The Ct value of real time PCR positive samples ranges between 19 and 34. A no-template control and positive control were included in every reaction (Fig. 5). The melting curve analysis of the PCR products showed typical melting profiles at 85°C (Fig. 6), while the negative samples did not show any melting curve.

Detection of *A. hydrophila* from different source

A total of 140 samples were collected from fish and different sources and subjected to culture on *A. hydrophila* selective media (AMB). From these, 81 (57.86%) isolates were presumptively identified as *A. hydrophila* by morphological and biochemical examination. These isolates were further confirmed as *A. hydrophila* by qPCR 54 (66.67%) positive as shown in (Fig. 4) based on specific primers on the sequence of the *ahaI* gene from the strain *A. hydrophila* subsp. *hydrophila* ATCC 7966 (Table 2).

Detection of *Aeromonas hydrophila* isolated from examined fishes based on the organs

In the current study, *A. hydrophila* was detected on the basis of their organ's location. Accordingly, the highest detection (40.54%) was assessed in both Muscle and gill, and the lowest (2.7%) was observed in Spleen (Table 3).

Antibiogram analysis

In the present study, antibiogram assay for the examined *A. hydrophila* isolates concerning 10 antibiotics revealed that all the tested isolates were completely sensitive to ciprofloxacin (100%), chloramphenicol (100%) and ceftriaxone (100%). In addition, amoxicillin and penicillin did not exhibit any bactericidal activity (100% resistant) as shown in (Table 4) and (Fig. 7) and multi antibiotic resistance index of 0.18 as shown in (Table 5). The result was interpreted as sensitive, intermediate and resistant according to the National Committee for Clinical Laboratory Standards (NCCLS) recommendations for *Aeromonas* species [15].

Discussion

Bacterial diseases are considered to be the most serious disease problem among freshwater fishes [16]. *A. hydrophila* has gained increased attention due to pathogenicity to humans and the ubiquity of the organism in the environment, food and water [17]. Isolation of *A. hydrophila* from four freshwater lake fishes along its value chain during the current study adds more evidence for the wide geographical distribution of the bacteria.

The clinical picture and postmortem findings observed in the current study of Nile tilapia were nearly similar to those described by [18–20]. The phenotypic and biochemical characteristics of *A. hydrophila* isolates recorded were in line to those reported in Bergey's manual of determinative bacteriology [21]. Similar phenotypic and biochemical findings with current study were also reported by [3, 18, 22–24].

Hemolytic activity of the isolates was determined for its importance as a virulent factor. Accordingly, from over all isolates 92.59% show β hemolysis and only 7.4% show α hemolysis. These toxins are responsible for lethality, hemolysis and entero-toxicogenicity. Their production




Fig. 4 Real time PCR positive samples of *A. hydrophila*




Fig. 5 Positive and Negative Controls




Fig. 7 Confirmed complete drug resistance pattern for P and AMP.

Table 5 Frequency distribution of multidrug resistant *A. hydrophila* isolates

Resistance pattern	<i>A. hydrophila</i> isolates (no = 10)		
	No. of <i>A. hydrophilaisolates</i>	Percentage of <i>A. hydrophilaisolates</i>	MAR index
Resistance to 2	10	100	0.28
Resistance to 3	6	60	0.2
Resistance to 4	2	20	0.08

Average MAR = 0.18

MAR: Multi - Antibiotic Resistance

(ahaI) constitutes an interesting and valuable study, not only to identify the specie, but also, enables future projects regarding recombinant adhesin as potential vaccine against Aeromonadaceae. From the total of 54 (66.67%) of qPCR positive samples, 37 (68.51%), 4 (7.4%), and 13 (24.07%) were from Fish source, RTE, and market fish respectively with no disease outbreak reported in all lakes at the point in time. As it was explained by Gilda [25], that disease occurrence in fish is a function of the pathogen, host and the environment. These results were at par with those reported by [26] in Iraq who found that over all detection rate of 65% *A. hydrophila* [27], in Berlin, Germany who found 63% cytotoxin producing *A. hydrophila*. However, lower prevalence were detected by [28] in Tamilnadu, India who found 40% of detection rate; [29] who found 40% of *A. hydrophila* from wild fish in Assiut, Egypt [23], in Moshtohor Egypt, who detected the total prevalence of bacterial infection (55.3%) [30], who found the prevalence of *A. hydrophila* 47% in Alexandria, Egypt, and [31] in Brazil who found the total prevalence of 46.66% *A. hydrophila*. However, a higher prevalence of *A. hydrophila* (95.06%) was reported by [32] in LiebefeM-Bern [33], in Kafrelsheikh governorate, Egypt who found a total prevalence of 75%. Variations

in the incidence level of *A. hydrophila* in the fish worldwide can be attributed to sampling time and geographical range [34]. Difference in the current study may be attributed to the number of examined fish, the size of fish and environmental conditions, geographical range, seasons of the study, sensitivity, and specificity of the techniques used to identify the bacteria.

Overall *A. hydrophila* (24.07%) contaminations in the market fish and RTE (7.4%) was observed in the current study. These results are in accordance with [35], who identified *A. hydrophila* (22.6%) from market fish in Ankara (Turkey). In Brazil [36] who detected 22.9% *A. hydrophila* from market fish samples. However, lower prevalence was detected by [37] Santos et al. (2002) who isolated 13% *A. hydrophila* from market fish samples in Brazil. Different studies have reported inconsistent detection rates of *A. hydrophila* for instance, Minana identified 2% of market fish in Spain, While, in India, 15.6% detection rate of *A. hydrophila* was reported in marketed fish samples by [38]. However, a higher prevalence recorded by [39] in Riyadh, Saudi Arabia who found 34% from fish market samples [29], who found (40%) of *A. hydrophila* from market fish in Assiut, Egypt and Attia [40], who reported overall higher *A. hydrophila* (51.4%) contaminations in the market fish in Sharkia Governorate, Egypt. This may be due to post-harvest contamination during selling through fishermen improper handling and transportation from the catching area. Fish in retail in the current study area are considered potential source for infection of human consumers. Although, the source of the organism may be ambient environment, secondary contamination during catching, handling and transportation may also contribute for its distribution.

Fish products ("leb-leb", fish salads, "gulash", smoked fish, etc.) are some of the most popular RTE choices in

Ethiopia. Concerning the detection of *A. hydrophila* in RTE fish, the current study revealed 7.4%. This result is in accordance with [40], who detected the prevalence of *A. hydrophila* in RTE grilled fish 8.6%. Mohamed [41], in Assiut Egypt, reported that *A. hydrophila* 20 and 10%, detection rate in grilled and fried fish samples respectively. A lower percentage (2.3%) of *A. hydrophila* was reported in RTE fish product in India by Gupta. Whereas, a higher percentage (77.3%) in RTE fried fish in India was also reported by [42]. The contamination rate in RTE fish may suggest contamination after cooking caused by lack of hygiene, contaminated water or contaminants from uncooked produce. The presence of *A. hydrophila* in RTE products again may be attributed to rapid grilling which could be insufficient to kill *A. hydrophila* that may be present in raw fish before preparation.

Regarding the frequency of detecting *A. hydrophila* from the different parts of the fish, out of 37 (68.51%) fish tissue samples, it was noticed that the highest (40.54%) gene detection was recorded from both gill and muscle respectively, (10.81%) from intestine, 5.4% from kidney and the lowest (2.7%) gene detection was recorded from spleen. The high proportion of infection in gills and muscle in comparison to other organs is due to the exposed nature of the organ to microbiota. The current findings are supported by the observations of [32, 43–45] who reported that *A. hydrophila* has been detected from wild fish, pond cultured edible and ornamental fish from different parts of the fish. These are attributed to the ubiquitous nature of the microorganism in the aquatic environment. The predominance of *A. hydrophila* in the gill and muscle of fishes may be attributed to the presence of *A. Hydrophila* in contaminated water in which the fish lives [5].

With the steady expansion of the fishery industry, the vast use of antibiotics will be unavoidable. The continuous and extensive use of antibiotics in humans also led to the emergence of antimicrobial-resistant strains worldwide [46]. Ten antibiotics namely; Ampicillin, Penicillin, Tetracycline, Ciprofloxacin, Chloramphenicol, Streptomycin, Gentamicin, Ceftriaxone, Amoxicillin-clavulanate, and Trimethoprim-Sulphamethoxazole were used in the current study mainly due to their routine usage in veterinary and human medicine. Fish treatments are not practiced almost in all fishery and aquaculture sectors of Ethiopia but, Tetracycline is commonly applied for the treatment of bacteremia in fishery research centers of Ethiopia (observation).

In the present study, antibiogram assay for the examined *A. hydrophila* isolates concerning 10 antibiotics revealed that all the tested isolates were completely sensitive to ciprofloxacin (100%), chloramphenicol (100%) and ceftriaxone (100%). In addition, amoxicillin and penicillin did not exhibit any bactericidal activity (100% resistant) against the tested isolates. These results are nearly agreed

with those obtained by [10, 11, 16, 19, 30, 47–50]. Freshwater streams are usually receptors of many industrial, domestic and agricultural wastes, which could contain antimicrobial agents and antimicrobial-resistant bacteria [51, 52]. Due to diverse microbial population in such ecosystems freshwater environment provides favorable conditions for the spread of antimicrobial resistance. The resistance to penicillin in *A. hydrophila* is mainly attributed to β -lactamase production that is encoded in their chromosomes. The antibiotic resistance has a public health concern as it mainly results from the improper intensive use of antibiotics [10]. The aeromonads have been regarded as being universally resistant to penicillin [10], in the current study penicillin and ampicillin resistance were confirmed. In the present study the multi-drug resistant (MAR) of the *A. hydrophila* were 0.18 and these findings are in accordance with the previous study of [10, 11].

Conclusions

The present study provided first evidence of infections of fish and fish products with virulent *A. hydrophila* strains. The pathogen was isolated and identified in 81 samples. On phenotypic assessments 92.59% (n=75) of the isolate expressed virulence trait of β -hemolysis. Molecular characterization using real-time PCR revealed presence of the adhesin gene (*ahaI*) in 54 (66.67%) of the isolates. Meanwhile, antimicrobial susceptibility test on selected *A. hydrophila* strains revealed the presence of resistance to amoxicillin and penicillin. The phenotypic and genotypic analysis provided epidemiological evidence for dissemination of a virulent *A. hydrophila* strain among the fish population in rift valley lakes. The detection of the pathogen in hemopoietic organ of the sampled fish population is alarming for potential outbreaks. The identified *A. hydrophila* isolates carry virulence trait that aids in colonization, infection and pathogenicity with ability to resist antibiotics commonly used in human and veterinary medicine. *A. hydrophila* is a zoonotic emerging pathogen and fish in lakes and fish products from Lake Koka, Zeway, Langano and Hawassa are potential sources of infection for humans in the area.


Methods

Study area

The current study was conducted in selected Rift Valley Lakes of Ethiopia, Koka, Ziway, Langano, and Hawassa from November 2020 to June 2021 from lake fishes, market fish and Restaurants of respective areas.

Study population

The present study was conducted on the Nile tilapia (*Oreochromis niloticus*), fish host biometric data including (average total body length TL, 246 mm, 5 mm accuracy) and weighed (average total body weight of 282.3 g)



Map of the study area

collected from the lakes, different fish markets and restaurants at study areas. Nile Tilapia were selected because of the fish population density and the trends of consumption preferences in the area [1].

Study design

A total of 140 fish samples were collected from different sources with a cross-sectional study from November 2020 to May 2021 at Koka, Ziway, Langan and Hawassa Lakes. The lakes were selected because of the bulk of the fish catch that contributes to 79% of the total fish catch in the country [1]. Restaurants in respective areas based on the accessibility to public transport transit areas and presence of recreational activities around the lake.

Sampling procedure

Purposive sampling strategy was followed in selecting fishes i.e. fish with suggestive lesions (hemorrhages on the external surface, the base of pectoral and tail fin, ulcer on the skin, abdominal distention, unilateral or bilateral

exophthalmia, prolapsed anus, and fin rot) of *A. hydrophila* infection were picked for sampling. As per mentioned somewhere tissue samples (muscle, gill, intestine, spleen and kidney) were collected from those fish having suggestive lesions [18–20]. All the fishes were caught using gillnets with mesh size ranging from (10 to 14 cm) that were used for the exploratory fishing work at the lakes. Samples were carried in Autoclavable sterile plastic bag containing water from the lake where they were caught and transported alive to Batu fishery and other aquatic life research center laboratory and Hawassa University Biology Department laboratory for post mortem examination and were analyzed immediately.

Clinical examination

Sampled fish were subjected to the clinical examination of the gross external signs as described by [18, 53, 54]. Fish was killed by transecting the spinal cord behind the skull. Autopsy and examination of the internal organs were carried out according to the method described by [55].

The organs sampled was muscle, gill, intestine, kidney, and spleen for bacterial culture and molecular analysis. First, the external body surface of the fish was examined for the presence of lesions, the gills, tail, and fins was observed for visible signs of infection and samples from muscle and gill were taken aseptically. After opening the body, the internal organs were exposed with care not to puncture any part of the intestinal tract by using ventral approach. In the absence of any visible lesions samples of a kidney, spleen and Intestine was taken after searing the surface of the organs with a hot scalpel blade. 2gm of each specimen were aseptically taken into the falcon tube (50ml) containing 20ml of alkaline peptone water PH 8.5 (Oxoid, England) which were kept cool at 4°C.

All raw fish was purchased and collected from traditional markets and supermarkets from respective towns by purposive selection based on the amount of fish stock kept on specific seller, market availability and customer's choice store based on informal collection of data in the area. Each market fish sample was individually packed in a clear sterile polyethylene bag immediate after sampling while, RTE fish samples collected as take away order and bagged in sterile plastic bags. 2gm of each specimen from market fish and RTE added to falcon tube containing 20ml of peptone water and preserved in an icebox. All the specimens from fish, market and RTE were labeled and transferred to the laboratory under aseptic condition with a minimum of delay in Batu and Hawassa University laboratories and finally brought to National Animal Health Diagnostic and Investigation Center (NAHDIC) with Electrical cooler jugs (icebox) for further studies.

Bacteriological examination (phenotypic identification)

A standard operating protocol was used for isolation and identification of *A. hydrophila* from fish and water samples [30, 56]. Aseptically taken 2gm of each fish sample (muscle, gill, intestine, kidney, and spleen) was thoroughly mixed (vortexed) from 20ml of samples in alkaline peptone water which is used as enrichment and transport media as per the method described by [22]. The homogenates were incubated for 24 h at 35 °C [53]. A loop-full from each enriched homogenate was streaked on to Aeromonas Medium Base (Oxoid, England), for each 500 ml Aeromonas medium base, 1 vial of ampicillin selective supplement was used and incubated for 24 h at 35 °C, a single colony from each suspected isolate was picked up and re-streaked on a new plate of its perused selective culture media and re-incubated at the same conditions. Presumptive colony from Aeromonas medium base inoculated in to Brain heart infusion broth (Oxoid, England) and incubated for 18-24 h at 35°C, then loop-full from the broth cultured on Nutrient agar media and incubated for 24 h at 35°C, each pure colony from the nutrient agar

medium used as a stock culture for further biochemical identification [57].

A. hydrophila were identified biochemically to species level based on colonial characteristics (colony morphology and arrangement) and by using 14 chosen biochemical test including gram staining of the microorganisms, cytochrome oxidase, catalase, motility, sugar utilization, indole, methyl red test, hemolysis production, Voges-Proskauer test, DNase test, gas production from Glucose, acid production from Sucrose, Mannitol and Xylose. Then the phenotypic and biochemical characteristics of the isolates were characterized according to the guideline indicated in Bergey's manual on fish and other aquatic animal practical identification manual [22].

Phenotypic characterization of *A. hydrophila* virulence determinants

The collected isolates were examined for their hemolytic activity on 5% whole sheep blood agar medium and results was recorded after 24 h of incubation at 35 °C and checked for the type (α or β) of hemolytic activity.

Molecular Detection of *A. hydrophila*.

DNA extraction

Genomic DNA was extracted using the DNA extraction kit (DNeasy kit, Qiagen, Germany) following the manufacturer's instructions. Qiagen DNeasy DNA extraction protocol for bacterial cultures adapted from Qiagen DNeasy handbook, 2020. Briefly, 200 μ l of the sample suspension was incubated at 70 °C for 10 min after the addition of 20 μ l of proteinase K and 200 μ l (AL) Buffer or lysis buffer by vortexing. Then, 200 μ l of 100% ethanol was added to the lysate and mixed thoroughly by vortexing. Washing and centrifugation of the sample was performed following the manufacturer's recommendations. Then, nucleic acid was eluted with 200 μ l of elution buffer provided in the kit.

Real-time qPCR amplification

Real-time qPCR was performed using a thermo cycler for real-time PCR (Applied Bio systems - Model Real time -7500) and the marker used was Eva green Super mix (Bio-Rad, USA). The Amplification reactions were performed in a reaction mixture of 20 μ l volumes consisting of 1 μ l of each ahaI primer (F and R), 10 μ l of 10x master mix including buffer, MgCl₂, dNTPs, Evagreen and DNA polymerase, 6 μ L of RNase-free distilled water and 2 μ l of genomic DNA template. The PCR program consisted of an initial step at 50 °C for 2 min and 95 °C for 10 min, followed by 40 cycles of denaturation at 95 °C for 15s and annealing at 60 °C for 1 min. At the end of each cycle, a DNA melting curve of the amplified products was performed between 65 and 95 °C, 95°C for 15 s, 65°C for 1 min and 95°C for 15 s with an increase of 0.5 °C in a

Table 6 Primer sequences (5' to 3') used to amplify the gene ahal in *A. hydrophila*, yielding a 200 bp amplicon

Primer	Primers sequences (5'-3')	T _m (°C)	Reference
ahal Forward	5- GAGAAGGTGACCAACAAGAACA-3	57.8	(58)
ahal Reverse	5- GAGATGTCAGCCTGTAGAGCT-3	54.2	

stepwise manner to evaluate the melting temperature (T_m) and to check the random amplification of untargeted regions.

Primer design

Gene specific primers used here were previously described by [58] based on the sequence of the ahal gene from the strain *A. hydrophila subsp. hydrophila* ATCC 7966. Sequences are shown in Table 6.

Antibiogram analysis

A. hydrophila strains was subjected to antibiotic sensitivity test using the Kirby-Bauer disc diffusion method according to the National Committee for Clinical Laboratory Standards (NCCLS) recommendations for *Aeromonas* species [15]. *A. hydrophila* isolates was inoculated in TSB and incubated at 35°C for 16-20 h, the turbid broth was inoculated in Muller Hinton broth (Oxoid, CM0405), the turbidity was adjusted according to McFarland obesity tube No. 0.5. Isolates was streaked on Muller Hinton agar (Oxoid, CM0337) and disks were placed, incubation was done at 37°C overnight. The used antibiotics were Amoxicillin-clavulanate (AMC, 30 µg), penicillin (P, 10 µg), Ampicillin (AMP, 10 µg), Ceftriaxone (CRO, 30 µg), Gentamicin (CN, 10 µg), Streptomycin (S, 10 µg), Tetracycline (TE, 30 µg), Ciprofloxacin (CIP, 5 µg), Trimethoprim-Sulfamethoxazole (SXT, 25 µg) and Chloramphenicol (C, 30 µg). Antimicrobials are selected based on the importance and common use in preventing and treating diseases in both veterinary and human medicines. After a period of 24 h. incubation, the zones of inhibition were compared and measured according to the manufacturer's instruction [15]. The result was interpreted as sensitive, intermediate and resistant according to the reference values.

The formula below is used to calculate the Multiple Antibiotic Resistances (MAR index) of the present isolates against tested antibiotics.

$$\text{MAR index} = X / (Y \times Z).$$

Where; X–Total of antibiotic resistance case.

Y–Total of antibiotic used in the study.

Z–Total of isolates. When the use of antibiotics is seldom or of low dose use for animal treatment, the MAR value is usually equal to or less than 0.2. In contrast, the elevated rate of use or the high risk of exposure of

antibiotics for animal treatment will yield an MAR index value which is more than 0.2.

Data management and analysis

The collected data were entered into Excel spreadsheet (Microsoft® office excel 2016) spread sheets and descriptive statistics was used.

Acknowledgements

My special thanks go to Dr. Delessa Damena for his helpful discussion, encouragement, providing constructive comments in improving the contents of the proposal. The Biology Department of Hawassa University and Batu fishery and other aquatic life research center acknowledged for providing laboratory space for the analysis during field work. I thank Ashu Bantashi from Batu for his valuable help in collecting samples.

Authors' contributions

NKK and AKA conducted fish sampling and bacterial sampling activities. NKK, AAM, MT, AOB,TKW, and DSG conducted laboratory work. NKK drafted the manuscript. TRC and YGM revised the manuscript and final approval of the submitted version.

Funding

This work was partially supported by Animal Health Institute. The funding sources had no role in study design; in the collection, analysis and interpretation of data; in the writing of the report; and in the decision to submit the article for publication.

Data Availability

The datasets used and analyzed during the current study are available from the corresponding author on reasonable request.

Declarations

Ethics approval

Ethical clearance was obtained from the animal research ethical review committee of Addis Ababa University College of Veterinary Medicine and agriculture for collecting samples from fishes during this study under the animal welfare guidelines and all methods were carried out in accordance with relevant guidelines and regulations. The research is also done according to the ARRIVE guidelines.

Consent for publication

Not applicable.

Competing interests

The authors declare no competing interests.

Author details

¹Department of Fish Disease Research and Diagnostics, Animal Health Institute, Sebeta, Ethiopia

²Animal Health Institute, Sebeta, Ethiopia

³College of Veterinary Medicine and Agriculture Department of Clinical Studies, Addis Ababa University, Debrezeit, Ethiopia

⁴Department of Molecular Biology, Animal Health Institute, Sebeta, Ethiopia

⁵Department of General Bacteriology, Animal Health Institute, Sebeta, Ethiopia

Received: 27 July 2022 / Accepted: 2 August 2023

Published online: 12 August 2023

References

1. FAO. Baseline Report for Ethiopia by Smart Fish Programme of the Indian Ocean Commission, Fisheries Management (FAO component). FAO, Ctry Rev. 2014;11–6.
2. Nicholson P, Mon-on N, Jaemwimol P, Tattiyapong P. Coinfection of tilapia lake virus and *Aeromonas hydrophila* synergistically increased mortality and worsened the disease severity in tilapia (*Oreochromis spp.*). *Aquaculture* [Internet]. 2020;520(2020):734746. Available from: <https://doi.org/10.1016/j.aquaculture.2019.734746>
3. Abdel-Latif HMR. Natural co-infection of cultured Nile tilapia *Oreochromis niloticus* with *Aeromonas hydrophila* and *Gyrodactylus cichlidarum* experiencing high mortality during summer. *Aquac Res*. 2020;1(1):1–13.
4. Mansour A, Mahfouz NB, Husien MM. MOLECULAR IDENTIFICATION OF *Aeromonas hydrophila* STRAINS RECOVERED FROM KAFRELSHEIKH FISH FARMS. *Slovak Vet Res*. 2019;56(22):201–8.
5. Fowoyo PT, Achimugu F. Virulence of *Aeromonas hydrophila* isolated from Fresh Water Catfish. *J Biosci Med*. 2019;7(12):1–12.
6. Ma S, Dong Y, Wang N, Liu J, Lu C, Liu Y. Identification of a new effector – immunity pair of *Aeromonas hydrophila* type VI secretion system. *Vet Res* [Internet]. 2020;1–14. Available from: <https://doi.org/10.1186/s13567-020-00794-w>
7. Zhou H, Gai C, Ye G, An J, Liu K, Xu L, et al. *Aeromonas hydrophila*, an Emerging Causative Agent of Freshwater-Farmed Whiteleg shrimp *Litopenaeus vannamei*. *Microorganisms*. 2019;9(12):10–8.
8. Zhong C. Comprehensive Analysis reveals the evolution and pathogenicity of *Aeromonas*. Viewed from Both Single Isolated Species and Microbial Communities. 2019;4(5):1–19.
9. Wu C, Ko W-C. *Aeromonas* isolates from Fish and patients in Tainan City, Taiwan: genotypic and phenotypic characteristics. *Am Soc Microbiol*. 2019;85(21):1–12.
10. Aa L, Najiah M. *Aeromonas hydrophila*: Antimicrobial susceptibility and histopathology of Isolates from Diseased Catfish, *Clarias gariepinus* (Burchell). *J Aquac*. 2013;5(2):1–15.
11. Stratev D, Odeyemi OA. Antimicrobial resistance of *Aeromonas hydrophila* isolated from different food sources: A mini-review. *J Infect Public Health* [Internet]. 2016;9(5):535–44. Available from: <https://doi.org/10.1016/j.jiph.2015.10.006>
12. Nuru A, Molla B, Yimer E, Health E, Box PO, Ababa A. Occurrence and distribution of bacterial pathogens of fish in the southern gulf of Lake Tana, Bahir Dar, Ethiopia. *Livest Res Rural Dev*. 2012;24(8):7.
13. Yimer E. Preliminary Survey of Parasites and bacterial pathogens of Fish at Lake Ziway. *Ethiop jSci*. 2000;23(1):25–33.
14. Almaw G, Woldeyes A, Adugna M, Koran T, Fen- A, Wubete A. Outbreak of *Aeromonas hydrophila* associated with the parasitic infection *Ichthyophthirius multifiliis* in pond of african catfish (*Clarias gariepinus*) fingerlings at Sebeta. *Ethiop Vet J*. 2014;18(2):109–16.
15. CLSIM45. M100 Performance Standards for Antimicrobial. 2020. 48–52 p.
16. Fernandez-Bravo A, Figueras MJ. An update on the Genus *Aeromonas*: Taxonomy, Epidemiology, and pathogenicity. *Microorganisms*. 2020;129(8):3–6.
17. Abdelsalam M, Ewiss MAZ, Khalefa HS, Mahmoud MA, Elgendy MY, Abdel-Moneam DA. Coinfections of *Aeromonas spp.*, *Enterococcus faecalis*, and *Vibrio alginolyticus* isolated from farmed Nile tilapia and African catfish in Egypt, with an emphasis on poor water quality. *Microb Pathog* [Internet]. 2021;160(July):105213. Available from: <https://doi.org/10.1016/j.micpath.2021.105213>
18. Austin B, Austin DA. Bacterial fish pathogens. *Bacterial fish pathogens*. 2016. 124–56 p.
19. Ramadan H, Ibrahim N, Samir M, El-moaty AA, Gad T. *Aeromonas hydrophila* from marketed mullet (*Mugil cephalus*) in Egypt: PCR characterization of β -lactam resistance and virulence genes. *J Appl Microbiol*. 2018;124(1):1629–37.
20. Abd-el-malek MA, Huihua Z, Chunlei G, Guifang Y, An J et al. crossm. *Vet World* [Internet]. 2019;9(2):1–12. Available from: <https://doi.org/10.1016/j.jiph.2015.10.006>
21. Garrity GM, Holt JG. The Road Map to the Manual. *Bergey's Manual® Syst Bacteriol*. 2001;119–66.
22. Buller NB. Bacteria and fungi from fish and other aquatic animals: A practical identification manual. *Bacteria and fungi from fish and other aquatic animals: A practical identification manual*. 2014.
23. Mohammed A, Aseely E, Shaheen A, Medicine FV, Gawad EAA, Abbass A. Phenotypic and molecular characterization of bacterial pathogens isolated phenotypic and molecular characterization of bacterial pathogens isolated from diseased freshwater fishes. *Int J Fish Aquat Stud*. 2018;6(2):34–41.
24. Skwor T, Stringer S, Haggerty J, Johnson J, Duhr S, Johnson M, et al. Crossm prevalence of potentially pathogenic antibiotic-resistant *Aeromonas spp.* *Treat Urban Wastewater Effluents versus*. 2020;86(3):1–16.
25. Gilda D. Health management in aquaculture. 2001. 122–125 p.
26. Aun Z, Alzainy A. The occurrence, hemolytic, cytotoxic activity and antibiotic susceptibility of *Aeromonas hydrophila* isolated from Fish samples in Baghdad. *Iraqi J Vet Med*. 2011;35(2):123–35.
27. Ullmann D, Krause G, Knabner D, Weber H, Beutin L. Isolation and characterization of potentially human pathogenic, cytotoxin-producing *Aeromonas* strains from Retailed Seafood in Berlin. Germany. 2005;87:82–7.
28. Nagarajan Y, Ramaian B. Detection of virulence gene in *Aeromonas hydrophila* isolated from Fish samples using PCR technique. *Glob J Biotechnol Biochem*. 2008;4(1).
29. Abd-el-malek AM. Incidence and virulence characteristics of *Aeromonas spp.* in fish. *Vet World*. 2017;10(January):34–7.
30. Dahdouh B, Basha O, Khalil S, Tanekhy M. Molecular characterization, Antimicrobial susceptibility and salt tolerance of *Aeromonas hydrophila* from Fresh, Brackish and Marine fishes. *Alexandria J Vet Sci*. 2016;48(2):46–53.
31. Rodrigues MV, Falcone-dias MF, Juliana C, David GS, José R, Pessoa J. *Aeromonas hydrophila* in Nile tilapia (*Oreochromis niloticus*) from brazilian aquaculture: a public health problem. *Emergent Life Sci Res*. 2019;5(1):48–55.
32. Gobat P, Jemmi T. Distribution of mesophilic *Aeromonas* species in raw and ready-to-eat fish and meat products in Switzerland. 1993;211:117–20.
33. Aboyadak IM, GMA1 N, Goda AA, Aboelgalagel WH, Alnokrashy A. Molecular detection of *Aeromonas hydrophila* as the Main cause of outbreak in Tilapia Farms in Egypt. *J Aquac Mar Biol*. 2015;2(5):2–5.
34. Vivekanandhan G, Hatha AAM, Lakshmanaperumalsamy P. Prevalence of *Aeromonas hydrophila* in fish and prawns from the seafood market of Coimbatore, South India. *Food Microbiol*. 2005;22:133–7.
35. Yucel N, Aslim B, Beyatli Y. *AEROMONAS SPECIES ISOLATED FROM. RETAIL FISH*. 2004;28(2005):313–24.
36. Sousa JA, De, Silva-souza AT. Bacterial Community Associated with Fish and Water from. *BRAZILIAN Arch Biol Technol AN*. 2001;44(4):373–81.
37. Santos JA, Otero A, Gonza MN. PCR detection of potentially pathogenic aeromonads in raw and cold-smoked freshwater fish. 2002;154:675–80.
38. I.Seethalakshmi R. Fisheries and aquatic science. *J offsheries Aquat Sci*. 2008;3(2).
39. Alhazmi MI. Isolation of. *Alhazmi J AOAC Int*. 2015;98(4):2014–6.
40. Attia ASA, Khedr MHE, Zaki MS. Occurrence of potentially pathogenic *Aeromonas* Species isolated from raw and ready-to-eat Fish marketed in Sharkia Governorate, Egypt. *Zagazig Veterinary J*. 2018;46(2):154–9.
41. Mohamed GM. COMPARATIVE STUDY BETWEEN RAW, AND COOKED FISH SOLD IN ASSIUT CITY ON THE INCIDENCE OF SOME FOODBORNE PATHOGENS. 2012;(April):99–108.
42. Sundaram E, Kathiravan S, Manna A, Chinnaiha A, Vasantha V. Ellairaja Sundaram, † Shenbagavalli Kathiravan, † Abhijit Manna, † Amutha Chinnaiha, † and VairathevarSivasamy Vasantha *, † †. *Am Chem Soc*. 2019;4(1):4814–24.
43. John N, Hatha AAM. Prevalence, distribution and drug resistance of motile aeromonads in freshwater ornamental fishes. *Indian J Fish*. 2012;59(2):161–4.
44. Nahar S, Rahman MM, Ahmed GU. Isolation, identification, and characterization of *Aeromonas hydrophila* from juvenile farmed pangasius (*Pangasianodon hypophthalmus*). *Int J Fish Aquat Stud*. 2016;4(4):52–60.
45. Rahman M, Colque-navarro P, Kühn I, Huys G, Swings J, Möllby R. Identification and characterization of pathogenic *Aeromonas veronii* Biovar Sobria Associated with Epizootic Ulcerative Syndrome in Fish in Bangladesh. 2002;68(2):650–5.
46. Salem M, Zahran E, Saad R, Zaki V. Prevalence, molecular characterization, virulotyping, and antibiotic resistance of motile aeromonads isolated from Nile tilapia farms at northern Egypt. *Mansoura Vet Med J*. 2020;21(1):56–67.
47. Lim SR, Lee D, Park SY, Lee S, Kim HY, Lee M, et al. Wild nutria (*Myocastor coypus*) is a potential Reservoir of Carbapenem-Resistant and zoonotic *Aeromonas spp.* in Korea. *Microorganisms*. 2019;7(2):1–14.
48. Baron S, Granier SA, Larvor E, Jouy E, Cineux M, Wilhelm A, et al. *Aeromonas* Diversity and Antimicrobial susceptibility in Freshwater — an attempt to set generic epidemiological cut-off values. *Front Microbiol*. 2017;8(2):1–9.
49. Igbinosa IH, Okoh AI. The scientific WorldJOURNAL Antibiotic Susceptibility Profile of *Aeromonas* Species isolated from Wastewater Treatment Plant. *Sci World J*. 2012;25(2):1–6.
50. Zdanowicz M, Mudryk ZJ, Perli P. Abundance and antibiotic resistance of *Aeromonas* isolated from the water of three carp ponds. 2020;9–18.

51. Mahsa Ansari. Antibiotic susceptibility and resistance of *Aeromonas* spp. isolated from fish. *Afr J Microbiol Res*. 2011;5(31):5772–5.
52. Von Graevenitz A. The role of *Aeromonas* in diarrhea: a review. *Infection*. 2007;35(2):59–64.
53. Austin B. *Bacterial Fish Pathogens Diseases of Farmed and Wild Fish*. Springer-praxis books in aquatic and marine sciences. 2007. 594–598 p.
54. Austin B. Infectious disease in aquaculture: Prevention and control. *Infect Dis Aquac Prev Control*. 2012;1–530.
55. Wu C-J, Ko W-C, Lee N-Y, Su S-L, Li C-W. *Am Soc Microbiol*. 2019;85(21):1–12.
56. Janda JM, Abbott SL. The Genus *Aeromonas*: taxonomy. Pathogenicity and Infection. 2010;23(1):35–73.
57. Balcázar JL, De Blas I, Ruiz-Zarzuela I, Vendrell D, Dolores Evora M, Luis Múzquiz J. Growth inhibition of *Aeromonas* species by lactic acid bacteria isolated from salmonids. *Microb Ecol Health Dis*. 2006;18(1):61–3.
58. Gertrudes E, Lemos DM, Pilarski F. Development of an Absolute Quantitative Real-Time PCR (qPCR) for the Diagnosis of *Aeromonas hydrophila* Infections in Fish *Acta Scientific Microbiology* (ISSN: 2581–3226) Development of an Absolute Quantitative Real-Time PCR (qPCR) for the Diagnosis. *ACTA Sci Microbiol*. 2018;1(4).

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.