

RESEARCH ARTICLE

Open Access



Effects of dietary postbiotic and inulin on growth performance, IGF1 and GHR mRNA expression, faecal microbiota and volatile fatty acids in broilers

Karwan Yaseen Kareem^{1,5}, Teck Chwen Loh^{1,2*}, Hooi Ling Foo^{3,4*}, Henny Akit¹ and Anjas Asmara Samsudin¹

Abstract

Background: Postbiotics (metabolic products by lactic acid bacteria) and prebiotics have been established as substitute to antibiotics in order to enhance immunity and growth performance in broiler chickens. Nonetheless, insufficient information is available on the effects of postbiotics and prebiotics combination on growth performance, faecal microbiota, pH and volatile fatty acids (VFA), as well as liver insulin like growth factor 1 (IGF1) and growth hormone receptor (GHR) mRNA expressions in broiler chickens. The aim of this experiment was to evaluate the effects of different types of postbiotics with different levels of prebiotic (inulin) on broiler for those parameters.

Results: The results showed that birds fed T3: (0.3 % RI11 + 0.8 % Inulin), T4: (0.3 % RI11 + 1.0 % Inulin), and T6: (0.3 % RG14+ 1.0 % Inulin) had higher ($p < 0.05$) final body weight (BW) and total weight gain (WG) than other treatments. Birds fed T3 had lower feed conversion ratio (FCR) which was significantly different from those fed with negative control diet but was similar to other treatments. Postbiotic and inulin increased ($p < 0.05$) faecal lactic acid bacteria (LAB) and reduced ($p < 0.05$) *Enterobacteriaceae* count. Birds fed T4 and T6 had higher faecal acetic acid and propionic acid respectively, and both had higher total VFA and lactic acid bacteria but lower pH and *Enterobacteriaceae* (ENT) counts compared to other treatments. The liver of birds fed T4 and T6 had higher IGF1 expression compared to other treatments while T6 had higher GHR mRNA expression compared to other treatments.

Conclusions: Results indicate that the addition of postbiotics and inulin combinations had beneficial effects on total BW, feed efficiency, mucosa architecture and IGF1 and GHR mRNA expression in broiler chickens.

Keywords: Broilers, Inulin, Prebiotic, Postbiotic, Intestinal microbiota, IGF1, GHR, Volatile fatty acid

Background

Intestinal microbiota play a vital role in the nutritional, physiological, immunological, and protective functions of the host [1] and their composition and activities can be influenced by diet [2]. The efficacy of feeding sub-therapeutic levels of antibiotics to modulate gut

microbiota to enhance production performance of live-stock has been espoused [3]. Unfortunately, the usage of antibiotics as feed additives for long periods in poultry diets can lead to antibiotic resistance [4] and high residue levels in poultry products such as meat and egg [5, 6]. Antimicrobial resistance encoding genes may represent risk to both human and animal health if it is transferred to other formerly susceptible bacteria [7]. Since the quest for safer and healthier chicken meat has remarkably increased in recent time, the use of natural feed additives can produce antibiotic-free chicken and can also prevent food-borne diseases [8].

* Correspondence: tcloh@upm.edu.my; hlfoo@upm.edu.my

¹Department of Animal Science, Faculty of Agriculture, Universiti Putra Malaysia, 43400 UPM, Serdang, Selangor, Malaysia

³Department of Bioprocess Technology, Faculty of Biotechnology and Biomolecular Science, Universiti Putra Malaysia, 43400 UPM, Serdang, Selangor, Malaysia

Full list of author information is available at the end of the article



In recent years, several feed additives such as prebiotics, probiotics, symbiotics, postbiotics and the combination of postbiotics and prebiotics have been used as growth promoters to replace antibiotics [9–12]. The mode of action of these additives differs. Probiotics colonize the host digestive system, increase the natural microbiota and prevent the colonization of pathogenic organisms [10]. Despite their beneficial effects, most probiotics especially the plasmids probiotics have antibiotic resistance genes which can be transferred between organisms [13]. As a consequence, probiotic as a live bacteria might not be used anymore in the near future. As a substitute to probiotics, metabolite products synthesized from probiotic known as postbiotics could be used. It is believed that postbiotics have the probiotic effects without living cells [14–16]. Prebiotics are non-living fibrous feed additives which when added to feed are preferred by harmful microbes. Prebiotics control the growth of pathogens (i.e. *Escherichia coli* and *Salmonella*) and stimulate the growth of *Bifidobacteria* and *Lactobacilli* and consequently promoting the health and performance of animals [17, 18]. A typical example of prebiotics is inulin. Postbiotics and inulin combination inhibited reproduction of pathogenic bacteria such as *Listeria monocytogenes*, *Salmonella enterica*, *Escherichia coli* and Vancomycin Resistant *Enterococci* [19]. Furthermore, addition of metabolite combinations to the feed of broilers [14, 15], laying hens [20] and pigs [16] improved the growth performance, faecal lactic acid bacteria and villus height. Various studies have examined the effects of postbiotic and prebiotics on growth performance, intestinal microbial ecology and histomorphology of broilers. However, there is dearth of information on the use of postbiotics and prebiotics combination and their synergistic effects on growth performance, intestinal microbial ecology, faecal VFA and histomorphology.

Chicken IGF1 has been identified as a biological candidate gene responsible for body composition, growth, fat deposition and metabolic activities in chickens [21]. It has been reported that the IGF-I level, feeding level, and growth rate are concurrent [22]. The dependence of nutritional and growth hormones on hepatic IGF1 production has been demonstrated [23, 24]. The pituitary releases the growth hormone which stimulates the hepatic production of IGF1 through the actions of GH activated GH receptors. However, the overall nutritional status of the animal modulates the ability of hepatic tissue to respond to GH. The IGF1 level can be affected by factors and situations that affect primary processes and control the IGF1 production [25]. The GHR gene play vital role as a mediator of body size in bird [26, 27]. Since probiotics, prebiotics, antibiotics and postbiotics influenced growth performance in poultry, a relationship between the feed additives and genes related to

growth is anticipated. Thus, the aim of this work was to examine the effect of postbiotics and prebiotics on growth performance, IGF1 and GHR expression, intestinal microbial ecology, histomorphology and faecal VFA in broilers.

Methods

Postbiotics and inulin

The stock culture of *Lactobacillus plantarum* (*L. plantarum*) RG14 and *L. plantarum* RI11 were prepared at the Laboratory of Prebiotic and Probiotic Technology II at Institute of Bioscience, Universiti Putra Malaysia. The stock cultures were revived two times using de-Mann Rogosa Sharpe (MRS) broth and incubated at 30 °C for 48 and 24 h subsequently at static condition, followed by spread plate and incubation was performed in 48 h at 30 °C. A single colony was then picked and inoculated into 10 mL MRS broth and incubated for 24 h. It was followed by subculturing it into 10 mL MRS broth and incubated for 24 h at 30 °C. The culture was then ready to be used as an inoculum. An inoculum size of 1 % (v/v) was inoculated into the respective reconstituted media and incubated for 24 h at 30 °C at static condition. Centrifugation at 10,000 × g for 15 min was performed to separate the bacterial cell. The postbiotics were collected and kept at 4 °C [28] prior to feeding trials. The inulin (Frutafit IQ) was provided by Connell Bros. Company (Malaysia) Sdn. Bhd.

Animals and experimental design

Two hundred and eighty-eight day old chicks were purchased from a commercial hatchery. The broiler chickens were allocated into eight treatment groups. Each group had six replicates while each replicate had six birds. The treatment groups included basal diet (negative control), basal diet + neomycin and oxytetracycline (positive control), T1 = Basal diet + 0.3 % postbiotic RI11, T2 = Basal diet + 0.3 % postbiotic RG14, T3 = Basal diet + 0.3 % postbiotic RI11 + 0.8 % inulin, T4 = Basal diet + 0.3 % postbiotic RI11 + 1.0 % inulin, T5 = Basal diet + 0.3 % postbiotic RG14 + 0.8 % inulin, T6 = Basal diet + 0.3 %, postbiotic RG14 + 1.0 % inulin. Water and feed were offered ad libitum to the birds until 42 days of age. Starter and finisher diets (Tables 1 and 2) were offered from days 0 to 21 and days 22 to 42, respectively. The experimental animals received humane care as outlined and approved by Institutional Animal Care and Use Committee for the Care and Use of Animals for Scientific Purposes (Research Policy, Universiti Putra Malaysia).

Timing of sample collection

On weekly basis, BW and feed intake (FI) were recorded and WG and FCR were calculated. For sampling, 12 birds per treatment group were slaughtered at day 42.

Table 1 Composition and nutrient content of starter diets

Ingredients	Dietary treatment ^a							
	Negative control	Positive control	T1	T2	T3	T4	T5	T6
Corn	50.00	50.00	50.18	50.18	50.20	50.20	50.20	50.20
Soybean	29.380	29.375	30.94	30.94	29.995	30.00	29.995	30.00
Wheat pollard	6.895	6.895	4.645	4.645	4.820	4.490	4.820	4.490
CPO	3.400	3.400	3.680	3.680	3.590	3.665	3.590	3.665
Fish meal (55 %)	7.580	7.575	6.825	6.825	7.550	7.600	7.550	7.600
L-Lysine	0.25	0.25	0.25	0.25	0.25	0.25	0.25	0.25
DL-Methionine	0.20	0.20	0.15	0.15	0.20	0.20	0.20	0.20
Monocalcium phosphate ²¹	1.00	1.00	1.10	1.10	1.00	1.00	1.00	1.00
Calcium carbonate	0.68	0.68	1.0	1.0	0.68	0.68	0.68	0.68
Choline chloride	0.06	0.06	0.06	0.06	0.06	0.06	0.06	0.06
Salt	0.25	0.25	0.30	0.30	0.25	0.25	0.25	0.25
Mineral premix ^b	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10
Vitamin premix ^c	0.06	0.06	0.06	0.06	0.06	0.06	0.06	0.06
Antioxidant ^d	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
Toxin binder ^e	0.135	0.135	0.400	0.400	0.135	0.135	0.135	0.135
Antibiotic ^f		0.01						
postbiotic RI11			0.30		0.30	0.30		
postbiotic RG14				0.30			0.30	0.30
Inulin					0.80	1.00	0.80	1.00
Calculated nutrient content (g/kg) ^g								
Crude protein	220.5	220.4	220.0	220.0	220.4	220.3	220.4	220.3
Metabolizable energy (MJ/Kg)	12.96	12.96	12.96	12.96	12.95	12.95	12.95	12.95
Calcium	9.9	9.9	10.01	10.01	9.9	9.9	9.9	9.9
Total phosphorus	8.5	8.5	8.3	8.3	8.5	8.3	8.3	8.3
Avail. P for poultry	5.1	5.1	5.1	5.1	5.1	5.1	5.1	5.1

^aNegative control: (basal diet), Positive control: (basal diet + neomycin and oxytetracycline), T1: (0.3 % RI11), T2: (0.3 % RG14), T3: (0.3 % RI11 + 0.8 % Inulin), T4: (0.3 % RI11 + 1.0 % Inulin), T5: (0.3 % RG14 + 0.8 % Inulin), T6: (0.3 % RG14 + 1.0 % Inulin)

^bMineral mix contains Fe 100 mg, Mn 110 mg, Cu 20 mg, Zn 100 mg, I 2 mg, Se 0.2 mg, Co 0.6 mg

^cVitamin premix contains retinol 2 mg, cholecalciferol 0.03 mg, α -tocopherol 0.02 mg, menadione 1.33 mg, cobalamin 0.03 mg, thiamine 0.83 mg, riboflavin 2 mg, folic acid 0.33 mg, biotin 0.03 mg, pantothenic acid 3.75 mg, niacin 23.3 mg, pyridoxine 1.33 mg

^dAntioxidant contains butylated hydroxyanisole (BHA)

^eToxin binder contains natural hydrated sodium calcium aluminium silicates

^fA combination of oxytetracycline and neomycin at the concentration of 100 ppm (w/w)

^gThe diets were formulated using FeedLIVE International software (Thailand)

FCR was calculated as follow: $FCR = \text{total feed consumed by birds} / \text{total weight gain}$.

Faecal LAB, ENT count and pH determination

The method of Foo et al. [29] was used to determine the faecal LAB and population of ENT. Faecal samples were kept at room temperature for 1 h once the 10-fold dilution (w/v) was done in sterile peptone water. Furthermore, 10-fold serial dilutions (v/v) were done after the 1 h soaking time. MRS-agar (Lactobacillus-Agar MRS) (Merck, KgaA, Darmstadt) was used to perform the enumerations of LAB. Incubation of

plates was performed in anaerobic jars for 48 h at 30 °C. The incubation of ENT was performed aerobically for 24 h at 37 °C after spreading and counting them on EMB-Agar (Eosin-methyleneblue Lactose Sucrose Agar, Merck, KgaA, and Darmstadt). The base 10 logarithm of colony-forming unit (CFU) (logCFU) per g was applied to express the number of CFU. The whole samples were in triplicates. Almost 9 ml of deionized distilled water was used to homogenise about 1 g of the sample in a universal tube. Mettler-Toledo pH meter with a glass electrode (Mettler-Toledo LTD, England) was used to measure the pH. The

Table 2 Composition and nutrient content of finisher diets

Ingredients	Dietary treatment ^a							
	Negative control	Positive control	T1	T2	T3	T4	T5	T6
Corn	54.70	54.70	54.89	54.89	54.80	54.69	54.80	54.69
Soybean	29.10	29.10	27.04	27.04	29.30	29.31	29.30	29.31
Wheat pollard	5.36	5.35	5.90	5.90	3.41	3.16	3.41	3.16
CPO	3.460	3.460	3.400	3.400	3.74	3.815	3.74	3.815
Fish meal (55 %)	3.600	3.600	5.040	5.040	3.870	3.945	3.870	3.945
L-Lysine	0.25	0.25	0.25	0.25	0.25	0.25	0.25	0.25
DL-Methionine	0.20	0.20	0.20	0.20	0.20	0.20	0.20	0.20
Monocalcium phosphate ²¹	1.40	1.40	1.35	1.35	1.40	1.40	1.40	1.40
Calcium carbonate	1.30	1.30	1.00	1.00	1.30	1.30	1.30	1.30
Choline chloride	0.06	0.06	0.06	0.06	0.06	0.06	0.06	0.06
Salt	0.25	0.25	0.25	0.25	0.25	0.25	0.25	0.25
Mineral premix ^b	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10
Vitamin premix ^c	0.06	0.06	0.06	0.06	0.06	0.06	0.06	0.06
Antioxidant ^d	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
Toxin binder ^e	0.15	0.15	0.15	0.15	0.15	0.15	0.15	0.15
Antibiotic ^f		0.01						
postbiotic RI11			0.30		0.30	0.30		
postbiotic RG14				0.30			0.30	0.30
Inulin					0.80	1.00	0.80	1.00
Calculated nutrient content (g/kg) ^g								
Crude protein	199.1	199.1	199.1	199.1	199.0	199.1	199.0	199.1
Metabolizable energy (MJ/Kg)	13.00	13.00	13.00	13.00	13.00	13.00	13.00	13.00
Calcium	10.3	10.3	10.0	10.0	10.5	10.5	10.5	10.5
Total phosphorus	8.1	8.1	8.2	8.2	8.0	8.0	8.0	8.0
Avail. P for poultry	4.7	4.7	4.9	4.9	4.7	4.7	4.7	4.7

^aNegative control: (basal diet), Positive control: (basal diet + neomycin and oxytetracycline), T1: (0.3 % RI11), T2: (0.3 % RG14), T3: (0.3 % RI11 + 0.8 % Inulin), T4: (0.3 % RI11 + 1.0 % Inulin), T5: (0.3 % RG14 + 0.8 % Inulin), T6: (0.3 % RG14 + 1.0 % Inulin)

^bMineral mix contains Fe 100 mg, Mn 110 mg, Cu 20 mg, Zn 100 mg, I 2 mg, Se 0.2 mg, Co 0.6 mg

^cVitamin premix contains retinol 2 mg, cholecalciferol 0.03 mg, α -tocopherol 0.02 mg, menadione 1.33 mg, cobalamin 0.03 mg, thiamine 0.83 mg, riboflavin 2 mg, folic acid 0.33 mg, biotin 0.03 mg, panthothenic acid 3.75 mg, niacin 23.3 mg, pyridoxine 1.33 mg

^dAntioxidant contains butylated hydroxyanisole (BHA)

^eToxin binder contains natural hydrated sodium calcium aluminium silicates

^fA combination of oxytetracyclin and neomycin at the concentration of 100 ppm (w/w)

^gThe diets were formulated using FeedLIVE International software (Thailand)

meter was calibrated prior to measuring the pH of the samples by using buffer solutions (Merck, KgaA, Dramstadt) at pH 4 and 7.

Histomorphology

Specimens were taken from three different parts of jejunum, duodenum, and ileum. These specimens were taken from the following locations;

- (i) the middle part of the duodenal loop,
- (ii) midway between the end point of duodenal loop and Meckel's diverticulum (jejunum), and

- (iii) midway between the Meckel's diverticulum and the ileo-caecal junction (ileum), and fixed in 10 % neutral buffered formalin.

A tissue processing machine (Leica, Japan) was used to excise, dehydrate these specimens. Then they were embedded in paraffin wax. Each sample was cut into sections (4 mm). These sections were stained with haematoxylin and eosin, fixed on slides, and then mounted and examined under the light microscope. The tip of the villus to the villus-crypt junction area was measured as the villus height. Furthermore, the crypt

depth was defined as the depth of the invagination between two villi.

Determination of VFA

The modified method of Thanh et al. [14] was applied to determine the VFA concentration in the faeces. From each sample, 1 g faeces (stored at -20°C) were weighed. Then, 1 mL of 24 % metaphosphoric acid was added which was diluted in 1.5 M sulphuric acid (BDH Laboratories, Poole, UK). First, under room temperature, this mixture was stored overnight and then the mixture was centrifuged at $10\,000 \times g$ for 20 min at 4°C . The supernatant was kept in a 1.5-ml screw-capped vial (Kimble Glass Inc., USA). For the GLC analysis, the internal standard, 20 mM 4-methyl-valeric acid (Sigma Chemical Co., St. Louis, MO, USA) was consistently added to the supernatant to make up 10 mM and the mixture was stored at -20°C . The VFA was separated on a Quadrex 007 Series (Quadrex Corp., New Haven, CT 06525, USA). The bonded phase was a fused silica capillary column (15 m, 0.32 mm ID, 0.25 mm film thickness) with a 6890 N (Hewlett-Packard, Avondale, PA) equipped with a flame ionization detector. The carrier gas was purified liquid nitrogen with a flow rate of 60 mL/min. The temperature of the injector and detector was set at 230°C . The column temperature was set at 200°C . For identification of peaks, the commercial standards of 20 mM acetic, and 10 mM each of propionic, butyric, isobutyric, valeric, isovaleric and 4-methyl-valeric acids from Sigma were used as external standards.

Total RNA isolation and reverse transcriptase polymerase chain reaction (RT-PCR) analysis of hepatic IGF1 and GHR

Lysis buffer (Qiagen, USA) received 30 mg (per sample) of frozen liver and it was properly homogenized. Following the manufacturer's instructions, Qiagen one-step RT-PCR kit (Qiagen, USA) was used to perform the RT-PCR. Real-time RT qPCR analyses were done using QuantiTect Primer Assay (200) IGF1 (QT00621334), GHR (QT00601321) and B-actin (QT00600614). Thereafter, NanoDrop was used to evaluate the purity of RNA at 260/280 OD ratio and RNA integrity. Only highly purified samples ($\text{OD}_{260}/\text{OD}_{280} > 1.8$) were chosen for further manipulation. The master mix preparation was also conducted following the manufacturer's procedure. The total volume of the reaction was 25 μl for each gene of interest arranged as $2 \times$ QuantiFast SYBR Green RT-PCR Master Mix 12.5 μl , 1 μl of forward primer, 1 μl of the reverse primer, 0.25 μl QuantiFast RT Mix, 1 μl of RNA, and finally 9.25 μl of RNase-free water. The reaction was carried out in a Bio-Rad thermal cycler (MyCycler, Germany). The RT-PCR conditions were as follows: (1) reverse transcription, 30 min, 50°C , (2) initial PCR activation step, 15 min, 95°C , (3) 3-step cycling for 40 cycles, each cycle consisting of denaturation for 30 s at

94°C followed by annealing for 30 s at $52\text{--}57^{\circ}\text{C}$ and extension for 1 min at 72°C . The linearity of response was ensured and the saturation of the reaction was averted by optimizing the template concentration and the cycle number. For standardization of the expression data, the β -actin mRNA fragment was used as internal standard (house-keeping gene). The results were standardized to the levels obtained for the β -actin gene. It was done by taking the ratio of the value obtained for the gene of interest to that of β -actin and then relative to the control. $2^{-\Delta\Delta\text{Ct}}$ ($\Delta\Delta\text{Ct} = \Delta\text{Ct Test sample} - \Delta\text{Ct Calibrator sample}$) calculated the relative mRNA expression

Statistical analysis

Data analysis was performed using the General Linear Model procedure of the Statistical Analysis System. Means were compared using the Duncan Multiple Range Test. The Bio-Rad CFX Manager 3.0 Software of the C1000 Touch thermal cycler-CFX96 Real time PCR (BIO-RAD, Foster city, California, USA) was used to calculate the relative gene expression of target genes in comparison to the β -actin reference gene.

Results

Growth performance

The growth performance of the birds fed diets containing different additives is presented in Table 3. Birds fed with T3, T4 and T6 had higher ($p < 0.05$) final BW and total WG than other treatments. The final BW and WG of birds fed the negative control diet, positive control diet, T1, T2 and T5 were similar ($p > 0.05$). There was no significant difference ($p > 0.05$) among the treatments for FI. Birds fed with T3 and T6 had lower ($p < 0.05$) FCR compared with birds fed the negative control diet. The FCR of birds fed T3, T4, T5, T6 and positive control were similar ($p > 0.05$). Similarly, the FCR of birds fed negative control, positive control, T1, T2, T4 and T5 diets did not differ.

Faecal LAB, ENT and pH

The faecal LAB, ENT and pH of birds fed various treatment groups are shown in Fig. 1. Significantly, the faecal pH for T3 were lower ($p < 0.05$) than the negative and positive controls. Dietary treatments affected ($p < 0.05$) LAB and ENT counts. Postbiotic and inulin increased ($p < 0.05$) faecal LAB and decreased ENT count when compared to negative control.

Histomorphology

The villus height and crypt depth of the duodenum, ileum and jejunum of birds fed different dietary treatments are shown in Table 4. Birds fed T3 and T6 had significantly higher ($p < 0.05$) villus height in the duodenum than the other treatments. Birds fed T2, T3, T4 and

Table 3 Growth performance at week 6 of treatments supplemented with different postbiotics and different levels of inulin

Parameter	Dietary treatments ^e								SEM
	Negative control	Positive control	T1	T2	T3	T4	T5	T6	
FI (g)	4245.17	4153.71	4289.19	4298.08	4199.29	4279.61	4187.49	4267.91	25.03
BW (g)	2239.59 ^b	2248.93 ^b	2267.28 ^b	2266.69 ^b	2334.90 ^a	2330.79 ^a	2264.72 ^b	2345.48 ^a	6.51
WG (g)	2189.10 ^b	2198.34 ^b	2217.79 ^b	2217.52 ^b	2284.31 ^a	2279.52 ^a	2215.59 ^b	2295.24 ^a	6.50
FCR	1.94 ^a	1.89 ^{abc}	1.93 ^{ab}	1.93 ^{ab}	1.84 ^c	1.88 ^{abc}	1.89 ^{abc}	1.86 ^{bc}	0.01

^{abc} means within a row for each parameter with different superscripts are significantly different ($p < 0.05$)

^enegative control: basal diet, positive control: basal diet + neomycin and oxytetracycline, T1: (0.3 % RI11), T2: (0.3 % RG14), T3: (0.3 % RI11 + 0.8 % inulin), T4: (0.3 % RI11 + 1.0 % inulin), T5: (0.3 % RG14 + 0.8%inulin), T6: (0.3 % RG14 + 1.0%inulin)

T6 had higher villus height in the ileum than the negative control. However, there were no significant differences ($p > 0.05$) for villus height in jejunum, and crypt depth in duodenum, jejunum and ileum among the treatments.

Volatile fatty acid

The faecal VFA of birds fed different dietary treatments is presented in Table 5. The result shows that acetic acid is the major VFA found in the broiler faeces followed by butyric and propionic acid. Broiler chickens fed T6 had the highest concentration of acetic acid and total VFA which was significantly different ($p < 0.05$) from other treatment groups. Birds fed T3 had the highest ($p < 0.05$) propionic acid as compare to birds fed with other treatments. No significant difference ($p > 0.05$) was observed for butyric acid in all treatment groups.

IGF1 and GHR mRNA expression

Gene expression profile of liver fed different dietary treatments is shown in Fig. 2. Birds fed T6 had the highest ($p < 0.05$) IGF1 mRNA expression. The IGF1 expression in T1, T3, T4, T5 and T6 were significantly higher ($p < 0.05$) than the negative and positive control. The GHR mRNA expression in the liver of broilers fed with

T6 was significantly higher ($p < 0.05$) than that of other treatment groups.

Discussion

Postbiotics influenced the growth rate of broiler chickens. It has bacteriostatic and bactericidal ability which decreases the pathogenic bacterial load in gastrointestinal microbiota. Kareem et al. [19] reported that *L. plantarum* exhibited inhibitory effect against various pathogens. Chicory root powder which comprise 68 % inulin enhances food digestion and absorption through jejunum histomorphometry modification thereby improve growth performance in broiler chickens [30]. The current results are in agreement with those of Thanh et al. [14] who found that birds fed combinations of metabolites produced by *L. plantarum* had higher ($p < 0.05$) final BW and WG compared with those fed the negative control diet. In contrast, Rosyidah et al. [31] did not observe significant difference in BW and WG in broilers fed with metabolites and combination of metabolite and acidifier groups and those fed positive and negative control diets.

The similarity in the BW and WG of birds fed the negative control diet and positive control diet suggests that antibiotics are no longer effective as growth enhancer and may not be effective as antibacterial in the future. The




Fig. 1 Microbiota counts (log CFU/g) and pH of broiler supplemented with different treatment diets at week 6. Treatments: Negative control: basal diet, Positive control: basal diet + neomycin and oxytetracycline, T1: (0.3 % RI11), T2: (0.3 % RG14), T3: (0.3 % RI11 + 0.8 % inulin), T4: (0.3 % RI11 + 1.0 % inulin), T5: (0.3 % RG14 + 0.8 % inulin), T6: (0.3 % RG14 + 1.0 % inulin). Bars with no common letter differ significantly ($p < 0.05$)

Table 4 Villus height and crypt depth in small intestine of treatments supplemented with different postbiotics and different levels of inulin

Parameter	Dietary treatments ^e								SEM
	Negative control	Positive control	T1	T2	T3	T4	T5	T6	
Villi height week 6, μm									
Duodenum	1304.88 ^b	1304.75 ^b	1306.80 ^b	1308.55 ^b	1419.68 ^a	1311.18 ^b	1315.80 ^b	1395.38 ^a	9.08
Jejunum	837.75	875.34	887.85	894.15	942.60	911.90	845.60	943.07	8.36
Ileum	540.90 ^c	595.71 ^{ab}	562.90 ^{bc}	617.30 ^a	634.08 ^a	610.87 ^a	597.25 ^{ab}	622.18 ^a	7.29
Crypt depth week 6, μm									
Duodenum	168.12	165.52	177.85	166.00	174.95	143.17	170.22	173.72	3.78
Jejunum	132.75	139.20	143.07	127.85	139.12	128.85	120.72	134.85	3.34
Ileum	101.07	106.15	105.50	102.12	107.42	105.75	95.15	109.45	2.67

^{abcd} means within a row for each parameter with different superscripts are significantly different ($p < 0.05$)

^enegative control: basal diet, positive control: basal diet + neomycin and oxytetracycline, T1: (0.3 % RI11), T2: (0.3 % RG14), T3: (0.3 % RI11 + 0.8 % inulin), T4: (0.3 % RI11 + 1.0 % inulin), T5: (0.3 % RG14 + 0.8%inulin), T6: (0.3 % RG14 + 1.0%inulin)

current observation is in tandem with that of Aristides et al. [32] who observed that dietary supplementation of avilamycin did not affect WG in broiler chickens. Postbiotics added with inulin was even better than diet added with antibiotic in BW and WG.

The lower FCR in T3 and T6 birds compared with the negative control birds is consistent with the report of Liu et al. [33] who found that probiotic, prebiotic and synbiotic significantly improved feed efficiency as compare to the negative control diet. In contrast, Elrayeh and Yildiz [34] reported 0.7 % inulin had no effect on FCR of broiler chickens.

Different combinations of postbiotics and inulin decreased faecal pH and ENT count. This observation corroborates the report of earlier findings which showed that addition of prebiotics beneficially modified intestinal microbiota in animal models and human studies. The authors posited that addition of prebiotics enhanced the population of protective bacteria (i.e. *Lactobacilli* and *Bifidobacteria*) and hindered the attachment of pathogenic bacteria to the gut epithelium [35–37]. The current findings are in agreement with those of Loh et al. [10] who observed that dietary postbiotics increased the faecal LAB and reduced the faecal pH and faecal ENT in laying hens. Also, Rosyidah et al. [31] observed an increase in LAB count and a decrease in ENT count

in the faeces of broilers fed with a metabolite produced by *L. plantarum*. Nabizadeh [38] found that dietary 1 % inulin reduced *E. coli* counts and pH in cecal contents in broiler chickens. Contrarily, Biggs et al. [39] found that corn-soybean meal diets containing 4 % inulin and some oligosaccharides had no effect on cecal *Lactobacillus*, *Bifidobacterium*, *Clostridium perfringens*, or *E. coli* populations in 21-d-old chicks. Abdel-Raheem et al. [40] also reported that prebiotic, probiotic and synbiotic had no effect on cecal *Lactobacilli* and *E. coli* counts in broilers chickens.

Villus height and crypt depth are reliable indicators of the gut function and health. According to Uni et al. [41] the health status of the gastrointestinal tract of an animal is a true reflection of intestinal morphology. Birds fed T3 and T6 had higher ($p < 0.05$) duodenal villus height than the control diets while birds fed T2, T3, T6 and T4 had higher ($p < 0.05$) ileal villus compared with the control diets. Our findings are consistent with the findings of earlier studies. The postbiotic and prebiotic altered the mucosal architecture in terms of longer villi and increased birds' performance [14, 40, 42]. Furthermore, probiotics and synbiotics can enhance broiler performance by improving the intestinal morphology and microbial balance which are associated with suppressing intestinal pathogens such as *E. coli* *Campylobacter* and

Table 5 Faecal VFA in broiler chickens fed with postbiotics and different levels of inulin

Parameter (mM)	Dietary treatments ^e								SEM
	Negative control	Positive control	T1	T2	T3	T4	T5	T6	
Acetic	27.78 ^c	29.78 ^c	30.39 ^c	34.70 ^{bc}	40.22 ^b	31.47 ^{bc}	36.85 ^{bc}	49.30 ^a	1.57
Propionic	0.27 ^c	0.66 ^{bc}	0.89 ^{ab}	0.71 ^{bc}	1.30 ^a	0.84 ^{ab}	0.87 ^{ab}	0.66 ^{bc}	0.07
Butyric	0.21	0.22	0.39	0.38	0.40	0.28	0.43	0.31	0.02
Total	28.25 ^d	30.65 ^{cd}	31.67 ^{cd}	35.45 ^{bcd}	41.92 ^b	32.59 ^{cd}	38.15 ^{bc}	50.27 ^a	1.59

^{abcd} means within a row for each parameter with different superscripts are significantly different ($p < 0.05$). ^enegative control: basal diet, positive control: basal diet + neomycin and oxytetracycline, T1: (0.3 % RI11), T2: (0.3 % RG14), T3: (0.3 % RI11 + 0.8 % inulin), T4: (0.3 % RI11 + 1.0 % inulin), T5: (0.3 % RG14 + 0.8%inulin), T6: (0.3 % RG14 + 1.0%inulin)




Fig. 2 IGF1 and GHR mRNA expression in the liver of broiler chicken. Treatments: Negative control: basal diet, Positive control: basal diet + neomycin and oxytetracycline, T1: (0.3 % RI11), T2: (0.3 % RG14), T3: (0.3 % RI11 + 0.8 % inulin), T4: (0.3 % RI11 + 1.0 % inulin), T5: (0.3 % RG14 + 0.8 % inulin), T6: (0.3 % RG14 + 1.0 % inulin). Bars with no common letter differ significantly ($p < 0.05$)

Salmonella and at the same time increase nutrient digestibility [43, 44]. In contrast, according to Nabizadeh [38], inulin did not affect the morphology of duodenum, ileum and jejunum of broiler chickens.

The observed increase in the concentration of acetic acid in T6, the higher propionic acid in T3 and increased in total VFA in both T3 and T6 compared with other treatments could be responsible for the increased lactic acid bacteria and decreased ENT counts and pH observed in the treatments. This finding is consistent with those of Loh et al. [10] and Thu et al. [16]. Postbiotics originating from *Lactobacillus* include valuable compounds such as organic acids and bacteriocin which enhance the growth of lactic acid bacteria [10]. Moreover, Van der Wielen et al. [45] reported that during growth of broiler chickens, VFA are responsible for the reduction in numbers of ENT in the ceca. In addition, the increased VFA could be due to the inulin as it contains polysaccharides and oligosaccharides. Gebbink et al. [46] demonstrated that fructooligosaccharide can be used to assist and maintain the healthy gastrointestinal tract environment by increasing the colonization of *Bifidobacteria* or reducing the *E. coli* in the intestinal system. Digestive enzymes do not hydrolyse the fructooligosaccharide in the small intestine of monogastrics and thus, it enters into the colon intact. Colonic microbiota metabolize it completely. Gases, lactate, and short chain fatty acid (i.e. acetate, propionate, and butyrate) are the output of carbohydrate fermentation [38]. The main fermentative chamber in broiler chicken is the caecum and this contain the largest number of bacteria compared with other gastrointestinal tract segment. Hence, the microbiota has high ability to ferment the carbohydrates [47].

Dietary postbiotic and inulin influenced the expression of mRNA IGF1 and mRNA GHR in the liver. The

increase in IGF-I mRNA and GHR mRNA in birds fed T6 is a true reflection of the growth performance of the birds. It is important to note that the IGF1 level, feeding level, and growth rate are concurrent [22]. The dependence of nutritional and growth hormones on hepatic IGF1 production has been demonstrated [23, 25]. Moreover, Amongst the genes influencing growth, IGF1 has been demonstrated as an indicator of growth rate in chicken by several authors [48, 49]. The pituitary releases the growth hormone which stimulates the hepatic production of IGF1 through the actions of GH activated GH receptors. However, the overall nutritional status of the animal modulated the ability of hepatic tissue to respond to GH [25]. The IGF1 level can be affected by factors and situations that affect primary processes and control the IGF1 production. These results might provide bases for the development of IGF1 as a growth index. Results from this study corroborate the findings of Beckman et al. [50] who found significant and positive correlation between mean growth rates of juvenile Chinook salmon and mean plasma IGF1 levels. However, other studies have not demonstrated any relationship between IGF1 and growth [51, 52] which can lead to uncertainties about the consistency of IGF1 growth relationships.

Conclusion

The study demonstrated that addition of postbiotics and inulin had beneficial effects on total BW, feed efficiency, mucosa architecture and expression of IGF1 and GHR mRNA in the liver of broiler chickens. However, birds fed T4: (0.3 % RI11 + 1.0 % inulin) and T6: (0.3 % RG14 + 1.0 % inulin) had higher total BW and feed efficiency than the other treatments. The faecal pH and ENT was reduced while VFA and LAB was increased in birds fed T4 and T6 as compared to the other treatments. Thus, both

T4 and T6 could be used as a substitute for antibiotics in broiler diet to improve the growth and gut health in broiler chickens.

Abbreviations

BW, body weight; CFU, colony-forming unit; ENT, *Enterobacteriaceae*; FCR, feed conversion ratio; FI, feed intake; GHR, growth hormone receptor; IGF1, insulin like growth factor 1; *L. plantarum*, *Lactobacillus plantarum*; LAB, lactic acid bacteria; MRS, de-Mann Rogosa Sharpe; RT-PCR, reverse transcriptase polymerase chain reaction.; VFA, volatile fatty acid; WG, weight gain.

Acknowledgements

The Long-Term Research Grant Scheme (LRGS) from Ministry of Education Malaysia has supported this project.

Funding

This project was supported by Long-Term Research Grant Scheme (LRGS) - Food Security entitled 'Enhancing the competitiveness and sustainability of the poultry industry for food security' from the Ministry of Higher Education, Malaysia.

Availability of data and materials

All relevant data and materials are available in the main manuscript.

Authors' contributions

KYK managed the chickens, TCL, HLF, HA, SAA participated in the whole design of the study and performed the statistical analysis. KYK carried out laboratory analyses and KYK, TCL, HLF, HA, SAA contributed to the preparation of the manuscript. All authors read and approved the final manuscript.

Competing interests

The authors declare that they have no competing interests.

Consent for publication

Not applicable.

Ethics approval and consent to participate

It is confirmed that Institutional Animal Care and Use Committee (IACUC), Universiti Putra Malaysia approved this study.

Author details

¹Department of Animal Science, Faculty of Agriculture, Universiti Putra Malaysia, 43400 UPM, Serdang, Selangor, Malaysia. ²Institute of Tropical Agriculture, Universiti Putra Malaysia, 43400 UPM, Serdang, Selangor, Malaysia. ³Department of Bioprocess Technology, Faculty of Biotechnology and Biomolecular Science, Universiti Putra Malaysia, 43400 UPM, Serdang, Selangor, Malaysia. ⁴Institute of Bioscience, Universiti Putra Malaysia, 43400 UPM, Serdang, Selangor, Malaysia. ⁵Department of Animal Resource, University of salah al-Din, Erbil, Iraq.

Received: 14 October 2015 Accepted: 3 August 2016

Published online: 05 August 2016

References

- Vispo C, Karasov WH. The interaction of avian gut microbes and their host: an elusive symbiosis. In: Mackie RI, White BA, editors. Gastrointestinal microbiology. New York: Chapman & Hall; 1997. p. 116-55.
- Rehman HU, Vahjen W, Awad WA, Zentek J. Indigenous bacteria and bacterial metabolic products in the gastrointestinal tract of broiler chickens. Arch Anim Nutr. 2007;61(5):319-35.
- Bolu SA, Adeyemi KD, Sola-Ojo FE, Fabiyi AA, Adedeji AT, Oluayemi O, Babalola AB. Lower dietary inclusion of alphamune g may improve performance, hematology, serum biochemistry and histology of broiler chicken. Sustainable Agr Res. 2012;1(2):15.
- Shazali N, Foo HL, Loh TC, Choe DW, Abdul RR. Prevalence of antibiotic resistance in lactic acid bacteria isolated from the faeces of broiler chicken in Malaysia. Gut Pathog. 2014;6(1):1.
- Shareef AM, Jamel ZT, Yonis KM. Detection of antibiotic residues in stored poultry products. Iraq J Vet Sci. 2009;23(1):45-8.
- Olatoye IO, Ehinwomo AA. Oxytetracycline residues in edible tissues of cattle slaughtered in Akure, Nigeria. Nigerian Vet J. 2010; 31(2):93-102.
- Montagne L, Pluske JR, Hampson DJ. A review of inter-actions between dietary fiber and the intestinal mucosa, and their consequences on digestive health in young non-ruminant animals. Anim Feed Sci Tech. 2003;108:95-117.
- Kleter GA, Marvin HJ. Indicators of emerging hazards and risks to food safety. Food Chem Toxicol. 2009;47(5):1022-39.
- Hajati H, Rezaei M. The application of prebiotics in poultry production. Int J Poult Sci. 2010;9(3):298-304.
- Loh TC, Choe DW, Foo HL, Sazili AQ, Bejo MH. Effects of feeding different postbiotic metabolite combinations produced by *Lactobacillus plantarum* strains on egg quality and production performance, faecal parameters and plasma cholesterol in laying hens. BMC Vet Res. 2014;10(1):149.
- Kareem KY, Loh TC, Foo HL, Asmara SA, Akit H, Abdulla NR, Ooi MF. Carcass, meat and bone quality of broiler chickens fed with postbiotic and prebiotic combinations. Int J Probiotics Prebiotics. 2015;10(1):23-30.
- Rebolé A, Ortiz L, Rodríguez ML, Alzueta C, Treviño J, Velasco S. Effects of inulin and enzyme complex, individually or in combination, on growth performance, intestinal microflora, cecal fermentation characteristics, and jejunal histomorphology in broiler chickens fed a wheat-and barley-based diet. Poultry Sci. 2010;89(2):276-86.
- Marteau P, Shanahan F. Basic aspects and pharmacology of probiotics: an overview of pharmacokinetics, mechanisms of action and side-effects. Best Pract Res Clin Ga. 2003;17(5):725-40.
- Thanh N, Loh TC, Foo HL, Hair-Bejo M, Azhar B. Effects of feeding metabolite combinations produced by *Lactobacillus plantarum* on growth performance, faecal microbial population, small intestine villus height and faecal volatile fatty acids in broilers. Brit Poultry Sci. 2009;50(3):298-306.
- Loh TC, Thanh NT, Foo HL, Hair-Bejo M, Azhar BK. Feeding of different levels of metabolite combinations produced by *Lactobacillus plantarum* on growth performance, fecal microflora, volatile fatty acids and villi height in broilers. Anim Sci J. 2010;81(2):205-14.
- Thu T, Loh TC, Foo H, Yaakub H, Bejo M. Effects of liquid metabolite combinations produced by *Lactobacillus plantarum* on growth performance, faeces characteristics, intestinal morphology and diarrhoea incidence in postweaning piglets. Trop Anim Health Pro. 2011;43(1):69-75.
- Boguslawska-Tryk M, Piotrowska A, Burlikowska K. Dietary fructans and their potential beneficial influence on health and performance parameters in broiler chickens. J Cent Europ Agri. 2012; 13(2):272-91.
- Jung S, Houde R, Baurhoo B, Zhao X, Lee B. Effects of galacto-oligosaccharides and a *Bifidobacteria* lactis-based probiotic strain on the growth performance and fecal microflora of broiler chickens. Poultry Sci. 2008;87(9):1694-9.
- Kareem KY, Ling FH, Chwen LT, Foong OM, Asmara SA. Inhibitory activity of postbiotic produced by strains of *Lactobacillus plantarum* using reconstituted media supplemented with inulin. Gut Pathog. 2014;6(1):23.
- Choe D, Loh T, Foo H, Hair-Bejo M, Awis Q. Egg production, faecal pH and microbial population, small intestine morphology, and plasma and yolk cholesterol in laying hens given liquid metabolites produced by *Lactobacillus plantarum* strains. Brit Poultry Sci. 2012;53(1):106-15.
- Kadlec J, Hosnedlová B, Řehout V, Čitek J, Večerek L, Hanusová L. Insulin-like growth factor-I gene polymorphism and its association with growth and slaughter characteristics in broiler chickens. J Agrobiology. 2011;28(2):157-63.
- Beckman BR, Shimizu MG, Gadberry BA, Parkins PJ, Cooper KA. The effect of temperature change on the relations among plasma IGF-1, 41-kDa IGFBP, and growth rate in postsmolt coho salmon. Aquaculture. 2004;241:601-19.
- Moriyama S. Increased plasma insulin-like growth factor-I (IGF-I) following oral and intraperitoneal administration of growth hormone to rainbow trout, *Oncorhynchus mykiss*. Growth Regulat. 1995;5(3):164-7.
- Shamblott MJ, Cheng CM, Bolt D, Chen TT. Appearance of insulin-like growth factor mRNA in the liver and pyloric ceca of a teleost in response to exogenous growth hormone. P Natl Acad Sci. 1995;92(15):6943-6.
- Beckman BR. Perspectives on concordant and discordant relations between insulin-like growth factor 1 (IGF1) and growth in fishes. Gen comp endocr. 2011;170(2):233-52.
- Li Y, Yang X, Ni Y, Decuyper E, Buyse J, Everaert N, Grossmann R, Zhao R. Early-age feed restriction affects viability and gene expression of satellite cells isolated from the gastrocnemius muscle of broiler chicks. J Anim Sci Biotechnol. 2012;3(1):33.
- Zhou N, Lee WR, Abasht B. Messenger RNA sequencing and pathway analysis provide novel insights into the biological basis of chickens' feed efficiency. BMC genomics. 2015;16(1):195.

28. Foo H, Loh T, Lai P, Lim Y, Kufli C, Rusul G. Effects of adding *Lactobacillus plantarum* I-UL4 metabolites in drinking water of rats. *Pakistan J Nutr.* 2003; 2(5):283–8.
29. Foo H, Loh T, Law F, Lim Y, Kufli C, Rusul G. Effects of feeding *Lactobacillus plantarum* I-UL4 isolated from Malaysian Tempeh on growth performance, faecal flora and lactic acid bacteria and plasma cholesterol concentrations in postweaning rats. *Food Sci Biotech.* 2003;12(4):403–8.
30. Izadi H, Arshami J, Golian A, Raji MR. Effects of chicory root powder on growth performance and histomorphometry of jejunum in broiler chicks. *Vet Res.* 2013;4(3):169–74.
31. Rosyidah MR, Loh TC, Foo H, Cheng X, Bejo MH. Effects of feeding metabolites and acidifier on growth performance, faecal characteristics and microflora in broiler chickens. *J Anim and Vet Adv.* 2011;10(21):2758–64.
32. Aristides LG, Paião FG, Murate LS, Oba A, Shimokomaki M. The effects of biotic additives on growth performance and meat qualities in broiler chickens. *Int J Poult Sci.* 2012;11:599–604.
33. Liu X, Yan H, Lv L, Xu Q, Yin C, Zhang K, Wang P, Hu J. Growth performance and meat quality of broiler chickens supplemented with *Bacillus licheniformis* in drinking water. *Asian Austral J Anim.* 2012;25(5):682.
34. Errayeh AS, Yildiz G. Effects of inulin and beta-glucan supplementation in broiler diets on growth performance, serum cholesterol, intestinal length, and immune system. *Turk J Vet Anim Sci.* 2012;36(4):388–94.
35. Santos A, San Mauro M, Diaz DM. Prebiotics and their long-term influence on the microbial populations of the mouse bowel. *Food Microbiol.* 2006; 23(5):498–503.
36. Langlands S, Hopkins M, Coleman N, Cummings J. Prebiotic carbohydrates modify the mucosa associated microflora of the human large bowel. *Gut.* 2004;53(11):1610–6.
37. Langen LV, Mirjam A, Dieleman LA. Prebiotics in chronic intestinal inflammation. *Inflamm Bowel Dis.* 2009;15(3):454–62.
38. Nabizadeh A. The effect of inulin on broiler chicken intestinal microflora, gut morphology, and performance. *J Anim Feed Sci.* 2012;21:725–34.
39. Biggs P, Parsons C, Fahey G. The effects of several oligosaccharides on growth performance, nutrient digestibilities, and cecal microbial populations in young chicks. *Poultry Sci.* 2007;86(11):2327–36.
40. Abdel-Raheem SM, Abd-Allah SM, Hassanein KM. The effects of prebiotic, probiotic and synbiotic supplementation on intestinal microbial ecology and histomorphology of broiler chickens. *Ijavms.* 2012;6:277–89.
41. Uni Z, Noy Y, Sklan D. Posthatch changes in morphology and function of the small intestines in heavy-and light-strain chicks. *Poultry Sci.* 1995; 74(10):1622–9.
42. Yang Y, Jji P, Choct M. Effects of different dietary levels of mannanoligosaccharide on growth performance and gut development of broiler chickens. *Asian Aust J Anim.* 2007;20(7):1084.
43. Ashayerizadeh A, Dabiri N, Ashayerizadeh O, Mirzadeh K, Roshanfekar H, Mamooee M. Effect of dietary antibiotic, probiotic and prebiotic as growth promoters, on growth performance, carcass characteristics and hematological indices of broiler chickens. *Pakistan J Biol Sci.* 2009;12:52–7.
44. Li X, Piao X, Kim S, Liu P, Wang L, Shen Y, Jung S, Lee H. Effects of chito-oligosaccharide supplementation on performance, nutrient digestibility, and serum composition in broiler chickens. *Poultry sci.* 2007;86(6):1107–14.
45. Van der Wielen PW, Biesterveld S, Notermans S, Hofstra H, Urlings BA, van Knapen F. Role of volatile fatty acids in development of the cecal microbiota in broiler chickens during growth. *Appl Environ Microb.* 2000; 66(6):2536–40.
46. Gebbink G, Sutton A, Richert B, Patterson J, Nielsen J, Kelly D, Verstegen M, Williams B, Bosch M, Cobb M. Effects of addition of fructooligosaccharide (FOS) and sugar beet pulp to weaning pig diets on performance, microflora and intestinal health. *Access.* 1999;12:53–9.
47. Józefiak D, Kaczmarek S, Rutkowski A. A note on the effects of selected prebiotics on the performance and ileal microbiota of broiler chickens. *J Anim Feed Sci.* 2008;17:392–7.
48. Jones JJ, Clemmons DR. Insulin-like growth factors and their binding proteins: biological actions. *Endocr rev.* 1995;16(1):3–34.
49. Beccavin C, Chevalier B, Cogburn LA, Simon J, Duclos MJ. Insulin-like growth factors and body growth in chickens divergently selected for high or low growth rate. *J Endocrinology.* 2001;168(2):297–306.
50. Beckman BR, Larsen DA, Moriyama S, Lee-Pawlak B, Dickhoff WW. Insulin-like growth factor-I and environmental modulation of growth during smoltification of spring chinook salmon (*Oncorhynchus tshawytscha*). *Gen comp endoc.* 1998;109(3):325–35.
51. Silverstein JT, Shearer KD, Dickhoff WW, Plisetskaya EM. Effects of growth and fatness on sexual development of chinook salmon (*Oncorhynchus tshawytscha*) parr. *Can J Fish Aquat Sci.* 1998;55(11):2376–82.
52. Nankervis L, Matthews S, Appleford P. Effect of dietary non-protein energy source on growth, nutrient retention and circulating insulin-like growth factor I and triiodothyronine levels in juvenile barramundi, *Lates calcarifer*. *Aquaculture.* 2000;191(4):323–35.

Submit your next manuscript to BioMed Central and we will help you at every step:

- We accept pre-submission inquiries
- Our selector tool helps you to find the most relevant journal
- We provide round the clock customer support
- Convenient online submission
- Thorough peer review
- Inclusion in PubMed and all major indexing services
- Maximum visibility for your research

Submit your manuscript at
www.biomedcentral.com/submit

